

First description of NDM-1-positive *Klebsiella pneumoniae* in the Tunisian community

Amel Mhaya, Rahma Trabelsi, Fatima M'zali, Dominique Bégu, Slim Tounsi,
Radhouane Gdoura, Corinne Arpin

► **To cite this version:**

Amel Mhaya, Rahma Trabelsi, Fatima M'zali, Dominique Bégu, Slim Tounsi, et al.. First description of NDM-1-positive *Klebsiella pneumoniae* in the Tunisian community. *Journal of Global Antimicrobial Resistance*, Elsevier, 2020, 21, pp.49-50. 10.1016/j.jgar.2020.02.023 . hal-02565943

HAL Id: hal-02565943

<https://hal-cnrs.archives-ouvertes.fr/hal-02565943>

Submitted on 6 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Letter to the Editor

First description of NDM-1-positive *Klebsiella pneumoniae* in the Tunisian community


Sir,

Multidrug-resistant bacteria, especially carbapenemase-producing Enterobacteriaceae, are a major public health-threat worldwide. As part of a collaborative monitoring programme, our laboratory at the University of Bordeaux has received a collection of multidrug-resistant bacterial strains to further characterise their β -lactamase content. They were sent from private Tunisian diagnostic laboratories and were collected from community patients suffering from urinary tract infection.

In this context, multidrug-resistant isolate 18TA was collected in January 2018 in Sfax region from the urine of a 45-year old female with no previous hospitalisation during the preceding month and no history of recent foreign travel.

Strain 18TA had been initially classified as *Enterobacter* spp. by biochemical tests (API 10S gallery). Following matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF/MS) (Bruker Daltonics) and confirmation by PCR amplification and sequencing of 16S rDNA, strain 18TA was re-identified as *Klebsiella pneumoniae*. Multilocus sequence typing (MLST) (<http://bigsd.bpasteur.fr/klebsiella/klebsiella.html>) indicated that strain 18TA belonged to Sequence Type (ST), ST147.

Minimum inhibitory concentrations (MICs) of various antimicrobials were determined using a BD PhoenixTM 100 automated system (BD Diagnostic Systems, Le Pont-de-Claix, France) and the results were interpreted using BD EpiCenterTM software (BD Diagnostic Systems). The MICs for ciprofloxacin and colistin were also determined by the broth microdilution method according to European Committee on Antimicrobial Susceptibility Testing 2019 guidelines (<https://www.sfm-microbiologie.org/2019/05/06/casfm-eucast-2019-v2/>).

Strain 18TA was resistant to all tested β -lactams, including carbapenems (Table 1). The strain was also resistant to gentamicin, tobramycin, quinolones (nalidixic acid), fluoroquinolones (ciprofloxacin) and trimethoprim/sulfamethoxazole (SXT) and showed decreased susceptibility to tigecycline (MIC = 2 μ g/mL). It remained susceptible to amikacin, fosfomycin and colistin (Table 1). The imipenem/ethylene diamine tetra-acetic acid (EDTA) combined disk diffusion test was positive since the inhibition zone increased by ≥ 7 mm with the imipenem/EDTA disk compared with

the imipenem disk alone, suggesting the presence of a metallo- β -lactamase (MBL) [1]. In addition, the double-disk synergy test (between amoxicillin/clavulanic acid and broad-spectrum cephalosporins) showed the presence of an extended-spectrum β -lactamase (ESBL)-producing phenotype (data not shown).

The transferability of the β -lactam resistance determinant was assessed by conjugation assay using an azide-resistant (Az^R) mutant of *Escherichia coli* C600 as the recipient strain. Selection was performed on Mueller–Hinton agar plates supplemented with sodium azide (300 μ g/mL) and ertapenem (4 μ g/mL). A transfer frequency of ca. 10^{-4} transconjugants per donor was observed. Comparison of MICs between the transconjugant (Tc-18TA) and its recipient strain (C600 Az^R) showed increased resistance not only to the tested β -lactams but also to gentamicin, tobramycin and SXT (Table 1).

Total genomic DNA of strain 18TA was screened using different multiplex PCR amplifications for various β -lactamase genes (*bla*_{TEM-like}, *bla*_{SHV-like}, *bla*_{OXA-1-like}, *bla*_{CTX-M} groups 1, 2, 9, 18 and 25, *bla*_{OXA-48-like}, *bla*_{KPC} and *bla*_{GES} and the MBL genes *bla*_{VIM}, *bla*_{IMP} and *bla*_{NDM}) as described previously [2]. Amplification results following agarose gel electrophoresis analysis showed the presence of group 1 *bla*_{CTX-M} and *bla*_{NDM} genes together with *bla*_{TEM-like}, *bla*_{SHV-like} and *bla*_{OXA-1-like} genes. Except for the *bla*_{SHV} gene that was found in strain 18TA but not in the transconjugant Tc-18TA and that was attributed to the chromosomally-encoded species-specific enzyme of *K. pneumoniae*, the four other β -lactamases were also found in transconjugant Tc-18TA (Table 1). Amplification of the entire *bla* genes was performed and subsequent sequencing ((Custom DNA sequencing; Eurofins Genomics GmbH, Ebersberg, Germany) showed the presence of the narrow-spectrum β -lactamase genes *bla*_{TEM-1B} and *bla*_{OXA-1} associated with the *bla*_{CTX-M-15} ESBL gene and the *bla*_{NDM-1} MBL gene both in 18TA and Tc-18TA (Table 1). Furthermore, amplifications searching for *aac(3)-IIa* (gentamicin and tobramycin resistance) and *sul1* and *dfrA1* (sulfamethoxazole and trimethoprim resistance, respectively) were positive both in 18TA and Tc-18TA. These genes were also present in Kp3771, a ST147 NDM-1-producing *K. pneumoniae* strain recently recovered from a patient hospitalised in an intensive care unit of University Hospital Tahar Sfar in Tunisia [3].

NDM-1-positive *K. pneumoniae* strains have been previously described in Tunisia, but only from hospitalised patients [1–5]. The current study reports the first description of *K. pneumoniae* carrying the carbapenemase NDM-1 in the Tunisian community

Table 1
 β-Lactamase (*bla*) gene content and antimicrobial susceptibility of *Klebsiella pneumoniae* 18TA and its *Escherichia coli* (Ec) transconjugant Tc-18TA.

Strain	<i>bla</i> gene content	MIC (μg/mL) ^a															
		TZP	FEP	ATM	IPM	MEM	ETP	GEN	TOB	AMK	SXT	NAL	CIP	FOS	TGC	COL	
18TA	<i>bla</i> _{SHV-like} <i>bla</i> _{TEM-1B} <i>bla</i> _{OXA-1}	>64	>16	>16	>8	>8	>1	>4	>4	≤4	>4	>16	128	32	2	1	
Tc-18TA ^b	<i>bla</i> _{CTX-M-15} <i>bla</i> _{NDM-1} <i>bla</i> _{TEM-1B} <i>bla</i> _{OXA-1} <i>bla</i> _{CTX-M-15} <i>bla</i> _{NDM-1}	>64	>16	>16	4	4	>1	>4	>4	≤4	>4	≤4	0.0625	≤16	≤0.25	0.25	
Ec C600 Az ^R	– ^c	≤4	≤1	≤0.25	≤0.25	≤0.25	≤0.25	≤1	≤1	≤4	≤1	≤4	0.0625	≤16	≤0.25	0.25	

MIC, minimum inhibitory concentration; TZP, piperacillin/tazobactam; FEP, cefepime; ATM, aztreonam; IPM, imipenem; MEM, meropenem; ETP, ertapenem; GEN, gentamicin; TOB, tobramycin; AMK, amikacin; SXT, trimethoprim/sulfamethoxazole; NAL, nalidixic acid; CIP, ciprofloxacin; FOS, fosfomycin; TGC, tigecycline; COL, colistin; Az^R, azide-resistant.

^a MICs were determined by BD PhoenixTM automated system. Strain 18TA and the transconjugant Tc-18TA were resistant to all β-lactams tested in the panel; only MICs to TZP, FEP, ATM, IPM, MEM and ETP are given. MICs for CIP and COL were also determined by the broth microdilution method.

^b In transconjugant Tc-18TA, the β-lactamase genes transferred by conjugation are reported, as well as the co-transferred resistances which are indicated in bold characters.

^c Not present.

setting and confirms that the NDM-1-positive ST147 *K. pneumoniae* clone has become endemic in this country [5]. Detection of this clone is of concern, therefore implementation of adequate infection control measures and uninterrupted active surveillance programmes both in hospital and community settings are crucial to prevent the dissemination of the NDM-1 enzyme in this region.

Funding

None.

Competing interests

None declared.

Ethical approval

Not required.

Acknowledgments

The authors thank Sabine Aillerie and Nathalie Peyron for their technical contribution. The authors also thank Véronique Dubois for performing MIC testing using the BD PhoenixTM automated system. This work was supported by the University of Bordeaux and National Center of Scientific Research (France) and by the Ministry of Higher Education and Research Scientific (Tunisia).

References

- [1] Izdebski R, Bojarska K, Baraniak A, Literacka E, Herda M, Żabicka D, et al. NDM-1- or OXA-48-producing Enterobacteriaceae colonising Polish tourists following a terrorist attack in Tunis, March 2015. *Euro Surveill* 2015;20:21150. doi:http://dx.doi.org/10.2807/1560-7917.es2015.20.23.21150.
- [2] Maamar E, Ferjani S, Jendoubi A, Hammami S, Hamzaoui Z, Mayonnove-Coulange L, et al. High prevalence of gut microbiota colonization with broad-spectrum cephalosporin resistant Enterobacteriaceae in a Tunisian intensive care unit. *Front Microbiol* 2016;7:1859. doi:http://dx.doi.org/10.3389/fmicb.2016.01859.
- [3] Mansour W, Grami R, Jaidane N, Messaoudi A, Charfi K, Ben Romdhane L, et al. Epidemiology and whole-genome analysis of NDM-1-producing *Klebsiella pneumoniae* KP3771 from Tunisia. *Microb Drug Resist* 2019;25:644–51. doi: http://dx.doi.org/10.1089/mdr.2018.0204.
- [4] Ben Nasr A, Decré D, Compain F, Genel N, Barguelli F, Arlet G. Emergence of NDM-1 in association with OXA-48 in *Klebsiella pneumoniae* from Tunisia. *Antimicrob Agents Chemother* 2013;57:4089–90. doi:http://dx.doi.org/10.1128/AAC.00536-13.

- [5] Messaoudi A, Haenni M, Mansour W, Saras E, Bel Haj Khalifa A, Chaouch C, et al. ST147 NDM-1-producing *Klebsiella pneumoniae* spread in two Tunisian hospitals. *J Antimicrob Chemother* 2017;72:315–6. doi:http://dx.doi.org/10.1093/jac/dkw401.

Amel Mhaya^{a,b,1}

^aUniversity of Bordeaux, UMR CNRS 5234, 146 rue Léo Saignat, Bordeaux Cedex, 33076, France

^bBiotechnology Center of Sfax, Laboratory of Biopesticides, BP 1177 Sfax, 3018, Tunisia

Rahma Trabelsi²

Department of Life Sciences, Research Laboratory of Environmental Toxicology–Microbiology and Health (LR17ES06), Faculty of Sciences, University of Sfax, Sfax, Tunisia

Fatima M'Zali

University of Bordeaux, Aquitaine Microbiologie, Bordeaux, France

Dominique Bégu

University of Bordeaux, UMR CNRS 5234, 146 rue Léo Saignat, Bordeaux Cedex, 33076, France

Slim Tounsi

Biotechnology Center of Sfax, Laboratory of Biopesticides, BP 1177 Sfax, 3018, Tunisia

Radhouane Gdoura

Department of Life Sciences, Research Laboratory of Environmental Toxicology–Microbiology and Health (LR17ES06), Faculty of Sciences, University of Sfax, Sfax, Tunisia

Corinne Arpin*

University of Bordeaux, UMR CNRS 5234, 146 rue Léo Saignat, Bordeaux Cedex, 33076, France

¹These two authors contributed equally to this article.

²These two authors contributed equally to this article.

* Corresponding author.

Received 10 October 2019

Available online 10 March 2020