

Editorial: Partitioning the Effects of Urbanization on Biodiversity: Beyond Wildlife Behavioural Responses to a Multilevel Assessment of Community Changes in Taxonomic, Functional and Phylogenetic Diversity

Federico Morelli, Piotr Tryjanowski, Anders Møller, Madhusudan Katti, Jiri Reif

► **To cite this version:**

Federico Morelli, Piotr Tryjanowski, Anders Møller, Madhusudan Katti, Jiri Reif. Editorial: Partitioning the Effects of Urbanization on Biodiversity: Beyond Wildlife Behavioural Responses to a Multilevel Assessment of Community Changes in Taxonomic, Functional and Phylogenetic Diversity. *Frontiers in Ecology and Evolution*, 2020, 8, 10.3389/fevo.2020.00023 . hal-03024727

HAL Id: hal-03024727

<https://hal-cnrs.archives-ouvertes.fr/hal-03024727>

Submitted on 26 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Editorial: Partitioning the Effects of Urbanization on Biodiversity: Beyond Wildlife Behavioural Responses to a Multilevel Assessment of Community Changes in Taxonomic, Functional and Phylogenetic Diversity

Federico Morelli^{1*}, Piotr Tryjanowski², Anders Pape Møller³, Madhusudan Katti⁴ and Jiri Reif⁵

¹ Department of Applied Geoinformatics and Spatial Planning, Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Prague, Czechia, ² Institute of Zoology, Poznan University of Life Sciences, Poznan, Poland, ³ Ecologie Systématique Evolution, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, Orsay, France, ⁴ Department of Forestry and Environmental Resources, Chancellor's Faculty Excellence Program for Leadership in Public Science, North Carolina State University, Raleigh, NC, United States, ⁵ Faculty of Science, Institute for Environmental Studies, Charles University in Prague, Prague, Czechia

Keywords: Anthropocene, biodiversity, cities, conservation, urban adapters, urbanization process

OPEN ACCESS

Edited and reviewed by:

Mark A. Elgar,
The University of Melbourne, Australia

*Correspondence:

Federico Morelli
fmorellius@gmail.com

Specialty section:

This article was submitted to
Urban Ecology,
a section of the journal
Frontiers in Ecology and Evolution

Received: 22 November 2019

Accepted: 28 January 2020

Published: 14 February 2020

Citation:

Morelli F, Tryjanowski P, Møller AP, Katti M and Reif J (2020) Editorial: Partitioning the Effects of Urbanization on Biodiversity: Beyond Wildlife Behavioural Responses to a Multilevel Assessment of Community Changes in Taxonomic, Functional and Phylogenetic Diversity. *Front. Ecol. Evol.* 8:23. doi: 10.3389/fevo.2020.00023

Editorial on the Research Topic

Partitioning the Effects of Urbanization on Biodiversity: Beyond Wildlife Behavioural Responses to a Multilevel Assessment of Community Changes in Taxonomic, Functional and Phylogenetic Diversity

THE CHALLENGE OF BIODIVERSITY CONSERVATION IN THE ANTHROPOCENE

We are living in the Anthropocene, where the effects of urbanization on biodiversity are increasingly important for global conservation (Aronson et al., 2014; Pickett et al., 2016). Cities can be considered refuges for some species but more likely ecological traps for others, because urban spaces constitute poor habitats offering little opportunity for reproductive success despite their apparent suitability. In this special Frontiers issue, we investigate a broad range of characteristics of cities in order to get a better understanding of the underlying mechanisms driving species distribution and abundance, and also the consequences of such proximate mechanisms for gaining a better understanding of the effects of urbanization on biodiversity.

Synurbanization is commonly defined as the process that leads the occupation of wildlife from rural to urban habitats, which can lead to an increase in the relative diversity of urban species. Urban habitats present new ecological challenges for animals, potentially filtering some species (Aronson et al., 2016). However, the diversity of urban habitat in terms of species composition can change depending on many aspects such as geographic location, buildings density, weather characteristics of the cities, and seasonal changes in communities. In this Research Topic, we tried to explore some of these potential factors and mechanisms driving changes in the behavior and composition of bird and mammal species assemblages.

Suhonen and Jokimäki examined whether urbanization has effects on the bird species wintering occupancy, distribution and abundance-occupancy relationship, and if such patterns varied between winters, considering the size of towns and other aspects of the Finnish landscape. The authors highlighted how three bird species were the core species during all winters, independently of the size of town or geographical latitude. Additionally, Suhonen and Jokimäki showed that the species abundance-occupancy relationship pattern was relatively stable over winters and across the sizes of towns.

On a different spatial scale and going deeper into the effects of urbanization on the gut microbiome of urban birds, Phillips et al. tested whether urbanization of habitats affects gut microbial composition and diversity in White-crowned Sparrows (*Zonotrichia leucophrys*). The authors found evidence that the birds' gut microbiome tends to be more diverse in urban than in rural environments, probably due to a higher variability of land cover types in cities than in farmlands. This evidence supports the statement that anthropogenic changes can affect species assemblages at multiple levels.

An interesting message for urban planning comes from the study of Hensley et al. focusing on comparison of biological traits between bird assemblages in three cities in south-western United States and their regional species pools. In two of three cities, namely Fresno, California and Tucson, Arizona, urban bird assemblages showed overrepresentation of granivores and omnivores, whereas insectivores were underrepresented relative to the regional species pool. This was not the case for Phoenix, Arizona, the largest of the focal cities, where representation of biological traits did not differ between the urban bird assemblage and the regional species pool. These striking contrasts between cities can be explained by local habitat composition: whereas the two smaller cities do not sample some of the surrounding native habitats, Phoenix contains protected mountain areas within the urban matrix of the city. This example shows how the inclusion of native habitats within the city borders can improve environmental conditions for urban biodiversity. On the other hand, Hensley et al. found that Fresno showed stronger filtering effects on the avifauna with the urban bird assemblage showing lower similarity with the regional species pool compared to Tucson or Phoenix. This may be driven by differences in urban water use policies resulting in greater extent of lawns causing Fresno to diverge from the more arid natural surroundings (Katti et al., 2017); Tucson (and to a lesser extent, Phoenix) on the other hand has stronger water conservation measures resulting in more xeriscaped yards, reducing the contrast with the desert habitats of the larger landscape matrix around that city. These findings highlight a role for more intentional careful landscape design and planning in modulating the filtering effect of urbanization on regional biodiversity.

However, the effects of urbanization are also evident for groups other than birds, such as, for example, mammals. Focusing on a typical adaptive behavior, Stillfried et al., have explored the risk perception and escape-behavior adjustment to urbanization in wild boar *Sus scrofa*. The authors used GPS tracking technology to monitor 11 wild boars from the metropolitan area of Berlin, in Germany. These data made it

possible to predict hotspots of potential human-wildlife conflicts. Finally, the authors also highlighted the behavioral plasticity of this species, which was able to adjust its escape behavior to human-dominated environments. This fact supports the statement that animals living in urban habitats are characterized by their acceptance of close proximity to humans (Hediger, 1934; Hemmingsen, 1951). The shorter flight initiation distance when approached by humans, perhaps reflecting a broader reduction in fear, constitute integral parts of the syndrome that characterizes urban animals (Morelli et al., 2018; Sol et al., 2018).

Changes in behavior, affected by urbanization processes, are visible not only during daylight, but also at night. And this problem is addressed by Straka et al. studying bats. Bats occur in urban areas around the world and nocturnal animals such as bats could be restricted in their activity by artificial light (Hopkins et al., 2018). Hence, understanding the impact of artificial light at night on this strictly nocturnal taxon is important for effective bat conservation in urban areas. The novelty of this study was the analysis of bat activity and light pollution in connection with tree cover density. They found a mitigating effect of tree cover for high-flying bat species in areas with high number of LED streetlamps. The study also reveals the complexity of the connections between artificial light and vegetation. The scientists recommend that tree-rich areas should only be illuminated sparingly and that trees should be planted in heavily illuminated areas to compensate. This could be a substantial contribution to the protection of urban bats and possible other nocturnal animals which can bring more nature to our cities.

CONCLUSION

Many of the focal factors in the collection of studies presented in this special issue impact on realized interactions between urban fauna and people. Urban areas attract biodiversity due to abundant food resources and heterogeneous habitats (McKinney, 2002), increasing the potential for wildlife-human interactions and conflicts. Considering the perspectives of urban ecology under future scenarios of land use and climate change, we highlight the necessity to articulate urban ecology as a multidisciplinary approach to better focus and assess the complexity of urban ecosystems (Pickett et al., 2016). We hope that the collection of studies provided in this special issue can help facilitate an understanding that even if cities are inexorably changing nature, many species are also adapting to urban environments, and in turn, this response from nature is also changing our perception of city life.

Our broad goal for this Research Topic was to take a deeper look at the effects of urbanization on biodiversity, going beyond descriptions of patterns to examine some of the important ecological, behavioral, and evolutionary mechanisms that drive species occurrence and interactions, and in turn, the species composition of urban communities. The papers in this collection offer a good sample of the variety of tools and approaches being employed to measure and analyse the diverse and complex effects of urbanization on biodiversity. While urban evolutionary ecology remains a relatively young

field, researchers are beginning to conduct deeper analyses of the recurrent patterns observed in the diversity and distribution of many taxa, testing both the generality of such patterns across cities, and uncovering key drivers and mechanisms that help explain the observed patterns. We expect future research to continue this trend, moving from phenomenological to mechanistic understandings of species occurrence and community dynamics in the context of urbanization. We also expect more sophisticated multivariate analytical models and experimental approaches to address the complex interactions and feedbacks between human activities and biodiversity, with the goal of developing better approaches for managing urban ecosystems in ways that reduce conflict and help reconcile human development and biodiversity conservation. As cities are becoming the preferred home of humanity, ecological, and evolutionary research must continue to expand its scope

beyond “natural” ecosystems and into the novel mixtures of natural and anthropogenic habitats that characterize cities as social-ecological systems. We suggest that this kind of research is in urgent need during this century of rapid global change, climate destabilization, and the ongoing mass extinction crisis.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

FM and JR were financially supported by the Czech Science Foundation GACR (project number 18-16738S).

REFERENCES

- Aronson, M. F. J., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A., et al. (2014). A global analysis of the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers. *Proc. R. Soc. Lond. B Biol. Sci.* 281:20133330. doi: 10.1098/rspb.2013.3330
- Aronson, M. F. J., Nilon, C. H., Lepczyk, C. A., Parker, T. S., Warren, P. S., Cilliers, S. S., et al. (2016). Hierarchical filters determine community assembly of urban species pools. *Ecology* 97, 2952–2963. doi: 10.1002/ecy.1535
- Hediger, H. (1934). Zur Biologie und Psychologie der Flucht bei Tieren. *Biol. Zent. Bl.* 54, 21–40.
- Hemmingsen, A. (1951). The relation of shyness (flushing distance) to body size. *Spolia Zool. Musei Hauniensis* 11, 74–76.
- Hopkins, G. R., Gaston, K. J., Visser, M. E., Elgar, M. A., and Jones, T. M. (2018). Artificial light at night as a potential agent of evolutionary change across the urban-rural landscape. *Front. Ecol. Environ.* 16, 472–479. doi: 10.1002/fee.1828
- Katti, M., Jones, A., Özgöç Çağlar, D., Delcore, H., and Kar Gupta, K. (2017). The influence of structural conditions and cultural inertia on water usage and landscape decision-making in a California Metropolitan Area. *Sustainability* 9:1746. doi: 10.3390/su9101746
- McKinney, M. L. (2002). Urbanization, biodiversity, and conservation. *Bioscience* 52, 883–890. doi: 10.1641/0006-3568(2002)052[0883:UBAC]2.0.CO;2
- Morelli, F., Mikula, P., Benedetti, Y., Bussière, R., Jerzak, L., and Tryjanowski, P. (2018). Escape behaviour of birds in urban parks and cemeteries across Europe: evidence of behavioural adaptation to human activity. *Sci. Total Environ.* 631–632, 803–810. doi: 10.1016/j.scitotenv.2018.03.118
- Pickett, S. T. A., Cadenasso, M. L., Childers, D. L., McDonnell, M. J., and Zhou, W. (2016). Evolution and future of urban ecological science: ecology in, of, and for the city. *Ecosyst. Heal. Sustain.* 2:e01229. doi: 10.1002/ehs2.1229
- Sol, D., Maspons, J., Gonzalez-Voyer, A., Morales-Castilla, I., Garamszegi, L. Z., and Møller, A. P. (2018). Risk-taking behavior, urbanization and the pace of life in birds. *Behav. Ecol. Sociobiol.* 72:59. doi: 10.1007/s00265-018-2463-0

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Morelli, Tryjanowski, Møller, Katti and Reif. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.