



HAL
open science

Reconstructing the Mesozoic-early Cenozoic evolution of northern Philippines: clues from palaeomagnetic studies on the ophiolitic basement of the Central Cordillera

Karlo Queaño, Jason Ali, Manuel Pubellier, Graciano Yumul, Jr., Carla Dimalanta

► To cite this version:

Karlo Queaño, Jason Ali, Manuel Pubellier, Graciano Yumul, Jr., Carla Dimalanta. Reconstructing the Mesozoic-early Cenozoic evolution of northern Philippines: clues from palaeomagnetic studies on the ophiolitic basement of the Central Cordillera. *Geophysical Journal International*, Oxford University Press (OUP), 2009, 178 (3), pp.1317-1326. 10.1111/j.1365-246X.2009.04221.x . hal-03080907

HAL Id: hal-03080907

<https://hal-cnrs.archives-ouvertes.fr/hal-03080907>

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License

Reconstructing the Mesozoic–early Cenozoic evolution of northern Philippines: clues from palaeomagnetic studies on the ophiolitic basement of the Central Cordillera

Karlo L. Queaño,¹ Jason R. Ali,² Manuel Pubellier,³ Graciano P. Yumul, Jr.^{4,5} and Carla B. Dimalanta⁵

¹*School of Earth and Materials Science and Engineering, Mapua Institute of Technology, Manila, Philippines*

²*Department of Earth Sciences, The University of Hong Kong, Pokfulam Rd., Hong Kong*

³*CNRS and Laboratoire de Geologie, Ecole Normale Supérieure, 24 rue Lhomond, Paris, France*

⁴*Department of Science and Technology, Bicutan, Taguig, Philippines*

⁵*National Institute of Geological Sciences, University of the Philippines, Diliman, Quezon City, Philippines. E-mail: rwm@mails@yahoo.com*

Accepted 2009 April 21. Received 2009 April 21; in original form 2008 January 18

SUMMARY

The first reliable palaeomagnetic data from the Cretaceous to Eocene ophiolitic basement rocks in the Philippines are presented. A total of 12 drill core sites from five localities in the Central Cordillera in northern Luzon, Philippines were sampled. Eight drill core sites were from pillow basalts, and four were from diabase feeder dykes. Combining the characteristic remanent magnetization direction from these sites gives a mean *in situ* direction of Dec = 162.2°, Inc = −21.4° ($\alpha_{95} = 17.0^\circ$, $k = 21.1$) and a tilt-corrected direction of Dec = 159.3°, Inc = −12.5°, $\alpha_{95} = 6.0^\circ$, $k = 162.5$. Along with other lines of palaeomagnetic evidence, this clustering suggests that the magnetization is primary and that the ophiolitic basement rocks of the Central Cordillera formed at subequatorial latitudes ($6.3^\circ\text{N} \pm 3.1^\circ$). This information further suggests that the basement rocks of northern Luzon were formed close to where the island was during the early Cenozoic. These rocks could represent relicts of the proto-Philippine Sea Plate.

Key words: Paleomagnetism applied to tectonics; Asia.

1 INTRODUCTION

As in the other regions of the western Pacific, reconstructing the Mesozoic–early Cenozoic evolution of the Philippine arc is still problematic. The lack of any model showing the evolution of the region during this period is likely a result of the limited information base that is available. It is long recognized, however, that Mesozoic ophiolite and ophiolitic complexes exist in the Philippines, which, when carefully studied, could provide clues on the early evolution of the region. Already, geochemical studies (e.g. Yumul *et al.* 2003; Tamayo *et al.* 2004; Encarnacion 2004; Queaño 2006; Yumul *et al.* 2006) have been conducted in many of these ophiolite sequences and their associated sedimentary carapace, the results of which point essentially to formation in subduction-related geodynamic settings. The question, however, still remains: did these units originate from a single marginal basin or from several marginal basins? In addressing this point, it is crucial that information relating to the palaeolatitude formation of these ophiolites be obtained. No reliable palaeomagnetic data have been reported previously from these rocks to determine the position of the basin at the time of their formation. This work reports the first reliable palaeomagnetic result from the

ophiolitic basement rocks in Luzon, providing crucial information on the origin of these rocks and conversely, the early evolution of the Luzon arc. The data were obtained from the Cretaceous–Eocene (?) ophiolitic basement rocks (herein referred to as the Chico River pillow basalts and dykes) in the Central Cordillera in northern Luzon.

2 TECTONIC SETTING OF NORTHERN LUZON

The Central Cordillera in northern Luzon forms part of the Philippine Mobile Belt (PMB), an active deformational zone bounded by two oppositely dipping subduction zones namely the Manila–Negros Sulu–Cotobato Trench to the west and the East Luzon Trough–Philippine Trench to the east (Fig. 1). The western trench system is where the marginal basins (South China Sea, Sulu Sea and the Celebes Sea) of the Eurasian Plate are subducting (Rangin & Pubellier 1990; Ringenbach *et al.* 1993). In the central Philippines, convergence of the Eurasian Plate with the PMB is expressed as a continent (Palawan Microcontinental Block)–arc collision zone. It is to be noted that the Palawan Microcontinental




Figure 1. Tectonic setting of the study area.

Block is that fragment in southern China, which translated south-eastwards, following the opening of the South China Sea sometime during the Oligocene (Lewis & Hayes 1984; Rangin 1989; Pubellier *et al.* 1991; Quebral *et al.* 1996; Yumul *et al.* 2003 2008).

The Philippine Trench and the East Luzon Trough comprise the eastern trench system, where the West Philippine Basin, the oldest marginal basin of the Philippine Sea Plate, is subducting. Activity along the eastern trench system is generally considered to be young. Ozawa *et al.* (2004) suggest southern propagation of subduction along the Philippine Trench from about 8 Ma, based on the temporal and spatial distribution of volcanic rocks along the east-

ern Philippines. Similarly, activity along the East Luzon Trough is thought to be young, given the shallow west-dipping Benioff zone associated with this plate boundary (Bautista *et al.* 2001). The presence of an eastward verging thrust zone observed in Taiwan (north of Luzon) also suggests incipient convergence between the Benham Plateau (a large igneous province within the West Philippine Basin) and Luzon along the East Luzon Trough (e.g. Cardwell *et al.* 1980; Stephan *et al.* 1986; Rangin & Pubellier 1990). The left-lateral Philippine Fault Zone accommodates the oblique convergence between the Philippine Sea and Eurasian plates bounding the Philippine Archipelago (Fitch 1972; Barrier *et al.* 1991; Aurelio 2000).




Figure 2. Stratigraphy of the study area.

3 GEOLOGY OF THE CENTRAL CORDILLERA

The Central Cordillera is the longest mountain range in the Philippines (Fig. 2). Geological data from the southern (Baguio City) and central (Lepanto-Tabuk) portions of the range suggest that it is a magmatic arc, formed mainly in response to subduction along the Manila Trench since the early Miocene (Bellon & Yumul 2000). This magmatic arc is floored by a Cretaceous-Eocene (?) ophiolitic complex, referred to as the Pugo Formation in the southern Central Cordillera. Correlatives of this formation are the Chico River pillow basalts and dykes exposed in the central portion of the range. Cenozoic volcanoclastic and epiclastic (mostly occurring as turbidites) rocks as well as reefal limestones overlie the ophiolitic basement rocks. Fig. 2 summarizes the stratigraphy of the Central Cordillera.

It is worth noting that Florendo (1994) combined the units of Pugo Formation with gabbroic plutons, mafic sheeted dykes and volcano sedimentary cover exposed in Baguio City in the southern Central Cordillera and adopted the term 'Itogon Ophiolite'. This was first introduced by the United Nations Development Programme (UNDP) (1987). He then assigned a late Oligocene to early Miocene age based on the radiometric dating of high-level gabbro and the age of the overlying volcanic carapace (Halfway Creek Formation). However, studies conducted by other workers (e.g. Pena 1992; Yumul 1994; this work) in Baguio City indi-

cate the presence of lithologies different from those reported by Florendo (1994). These studies report diorite, quartz diorite, granodiorite and andesites as the dominant igneous bodies rather than the gabbros reported by Florendo (1994). They were emplaced in different pulses during the late Oligocene to Pliocene (Maletterre 1989; Bellon & Yumul 2000). Florendo (1994) also reported the sedimentary carapace (the Halfway Creek Formation) as being composed of a sequence of basaltic pillows, pillow breccia and conglomerates, turbiditic wackes and rare massive rhyolite flows, red shales, calcarenites, calcirudites and micritic limestone. However, field mapping of areas where the Halfway Creek Formation is supposedly exposed, reveals widespread exposures of tuffaceous conglomeratic sandstone (part of the Zigzag Formation) and limestone (Kennon Limestone; Pena 1992; this work). Only minor volcanic flows were encountered. Encarnacion (2004) also recognized that some of the areas mapped by Florendo (1994) as sheeted dykes were, in fact, fractured lava flows.

Although the findings of Florendo (1994) appear to be in conflict with those obtained by other workers, there seems to be a consensus suggesting that the Central Cordillera has an ophiolitic basement. For instance, Maletterre (1989) introduced the Pingkian Ophiolite for the gabbro-pillow-basalt-dyke complex association presumably of pre-Oligocene age in the southern Cordillera. Ringenbach (1992) also recognized the ophiolitic units (largely massive and pillow basalts and sheeted dykes) of the Cretaceous?

Lepanto Metavolcanics as forming the basement of Lepanto-Cervantes region situated in the central part of the range. Yumul *et al.* (1992) argued for a possible correlation between the Lepanto Metavolcanics and Pugo Formation and further suggested formation of these units in a subduction-related marginal basin based on the chemistry of the volcanic-hypabyssal rocks.

Obtaining a reliable radiometric age for the ophiolite basement of the Central Cordillera is unsuccessful to date (including this work). However, previous authors commonly agree on a Cretaceous, possibly as young as Eocene age, for the basement rock of the Central Cordillera, based on stratigraphic relationships with the overlying sediments.

4 PALAEOMAGNETIC SAMPLES AND TECHNIQUES

Palaeomagnetic samples were obtained mainly along the Chico River, between Bontoc and Tabuk, in the central portion of the Central Cordillera (Fig. 2). In that locality, good exposures of ophiolitic basement rocks exist, mostly of pillow lavas intruded locally by diabase dykes. The orientation of the pillow lavas varies between outcrops, but generally is towards the west. Attempts were also made to collect samples from the Pugo Formation in the southern Central Cordillera. Unfortunately, exposures are limited and are too weathered and/or fractured for palaeomagnetic sampling.

A total of 12 drill core sites from five localities (~9 km stretch of ground) were sampled. Eight drill core sites were from pillow basalts whereas four were from diabase feeder dykes. A portable gas powered rock-drill was used to obtain six to eight, 2.54-cm-diameter core samples at each site.

Stepwise alternating field and/or thermal demagnetization were applied in the core specimens. Demagnetization behaviour was examined by visual inspection of vector endpoint diagrams (Zijderveld 1967) and equal area stereographic projections. The characteristic and secondary magnetization components were identified via principal component analysis (Kirschvink 1980) and/or the use of the statistics of Fisher (1953). Remanence carriers were determined by conducting isothermal remanent magnetization (IRM) and the Lowrie test (1990). The former experiment also involved looking at the IRM ratio developed by Ali (1989, in Ali & Hall 1995; Ali *et al.* 2001), with a ratio above 0.9 being indicative of magnetite as the dominant carrier. Demagnetization spectra of the IRMs adopting Lowrie's method provided more concrete results, suggesting remanence contribution from different carriers.

Specimens displaying chaotic behaviour (i.e. those with components having a maximum angular deviation or MAD $\geq 10^\circ$) were discarded in the determination of the site mean directions. The ratio of natural remanent magnetization (NRM) to the IRM demagnetization technique developed by Fuller *et al.* (1988) and Cisowski *et al.* (1990) was also applied on representative specimens to discriminate primary remanence from secondary chemical signal. This method is particularly useful for specimens bearing fine magnetite in that primary thermoremanent magnetization and secondary chemical signal are suggested when NRM/IRM ratios are greater than 10^{-2} and less than 10^{-3} , respectively.

5 PALAEOMAGNETIC RESULTS

5.1 Pillow basalt sites

Most pillow basalt samples (e.g. LZ15, 16, 17 and 22) have low NRM intensities, with values ranging from 0.2 to 4 mAm⁻¹. Typi-

cally, the pillow basalt specimens have two component remanences comprising a low coercivity signal, usually removed at ~10 mT (e.g. LZ15, 16 and 17), and a higher stability characteristic remanent magnetization (ChRM; Fig. 3). Site LZ22 has a significant low coercivity component (LCC) removed at ~25 mT. The mean direction of the LCC in most sites is poorly defined ($\alpha_{95} > 15^\circ$). However, when the LCC directions of the samples (with MADs $< 10^\circ$) are plotted, most *in situ* directions cluster near to that of the present field, suggesting that the LCC is likely a VRM (viscous remanent magnetization).

The direction of the ChRM was determined using the statistics of Fisher (1953) instead of the conventional principal component analysis. The former method provided a more precise determination of the ChRM direction of specimens, especially those having weak magnetizations and displaying somewhat 'noisy' behaviour. Three sites (LZ11, 19 and 20) were excluded from palaeomagnetic-tectonic modelling due to secondary overprinting or inconsistency in demagnetization behaviour of the ChRM components. Four sites (LZ15, 16, 17 and 22) have ChRM directions that group well at the sample mean level ($\alpha_{95} < 10^\circ$; Fig. 4). In tilt-corrected coordinates, the latter sites have SSE declinations and negative shallow inclinations. It is noteworthy that a specimen from LZ15 has a direction antipodal to that of the other specimens. In contrast, site LZ14 yielded a marginally acceptable ChRM grouping ($\alpha_{95} = 3.9^\circ$). However, its inclination (+48.4°) is anomalously steep compared with the other pillow basalt sites, and hence, the direction from this site is excluded from tectonic modelling. The mean direction of the four pillow basalt sites with $\alpha_{95} < 10^\circ$ is: *in situ* Dec = 164.2°, Inc = -18.5°, $\alpha_{95} = 21.4^\circ$, $k = 19.4$; tilt-corrected Dec = 159.9°, Inc = -13.3°, $\alpha_{95} = 7.8^\circ$, $k = 139.7$. The significant improvement in the direction clustering statistics suggests that the remanence predates folding (Table 1).

The magnetic carriers in the samples were determined using IRM experiments. Most representative pillow basalt specimens from the different localities yielded IRM ratios of 0.99, which suggests that magnetite is the principal remanence carrier (Fig. 5; Table 1). In two of the sites, some remanence contribution from maghemite and titanomagnetite is deduced from the LCC and intermediate coercivity component (ICC) fractions and the discontinuity in the LCC and ICC curves (Figs 6 and 7). Nonetheless, all specimens were adequately demagnetized and the calculated ChRM directions are consistent between specimens from each site (particularly those with $\alpha_{95} < 10^\circ$). In addition, the application of tilt correction has also significantly improved the clustering statistics of the ChRM mean direction. Although maghemization may have resulted in a decrease in magnetic intensity, the process appears to have imparted negligible effects on the rocks, which allowed for the preservation of primary magnetization directions.

Very little palaeomagnetic information is available on palaeogene to Early Neogene rocks in Luzon. Nevertheless, Fuller *et al.* (1983) have reported Eocene rocks from the Zambales ophiolite and associated formations in nearby Tarlac and Pangasinan as characterized by shallow inclination directions (mean $I = -8^\circ$). Similarly, Queaño (2006) has also shown Upper Eocene clastic sedimentary rocks of the Bangui formation (mean $I = 2.9^\circ$) and Late Oligocene–Early Miocene Palaui pillow basalts (mean $I = 6.3^\circ$) to lie intermediate between the calculated inclination angle for the Cretaceous–Eocene Chico pillow basalts (mean $I = -12.5^\circ$) and the Middle Miocene Klondyke Formation (mean $I = 19.0^\circ$; Queaño 2006) or the present-day geomagnetic inclination in the region ($I = 19.667^\circ$; IGRF 2005). Therefore, all things considered, there is sufficient basis to assume that the isolated ChRM and calculated




Figure 3. Vector endpoint plots (Zijderveld 1967) showing examples of demagnetization data in tilt-corrected coordinates of ophiolitic basement rocks along the Chico River and adjacent areas. Solid/open symbols are projections onto the horizontal/vertical planes.

palaeoinclination directions are indeed primary and are representative of the magnetization acquired by the lavas at crystallization during the Cretaceous.

5.2 Dyke sites

Three (LZ12, 13 and 21) out of four sites from the diabasic dykes have directions that could be evaluated for palaeomagnetic–tectonic modelling (Site LZ18 was rejected as the specimens exhibit different/erratic demagnetization behaviour). The dykes likely act as feeders to the pillows (and hence, they are magnetically coeval). NRM intensities of the lava flow specimens are variable, with values ranging from 0.2–400 mA m⁻¹. Sites LZ12 and 13 yielded single components of magnetization (Fig. 4). Specimens from LZ12 fully demagnetized at 35–50 mT, whereas those from LZ13 have ~20 per cent of the initial NRM remaining at 100 mT. Subsequent thermal demagnetization of specimens from the latter site did not give any useful results, as the specimens exhibited erratic magnetization behaviour. In contrast, site LZ21 yielded two components: an LCC, removed at 10–23 mT, and a higher stability ChRM, which converges to the origin ≥13 mT. It is noted, however, that three specimens from LZ21 do not directly converge to the origin, and that they showed noisy and very weak (<0.3 mA m⁻¹) magnetizations at the last stages of demagnetization treatment. As such, it was not possible to isolate the ChRM of the specimens by the conventional principal component analysis. Instead, ChRM directions of the specimens were estimated using Halls’ (1978) converging remagnetization circles method.

ChRM directions for the dykes group well at the sample level with $\alpha_{95} = 15^\circ$. However, the directions do not show any common trend in both *in situ* and tilt-corrected (i.e. using that applied for

the pillow basalts assuming a coeval magnetization) coordinates. The ChRM of LZ13 appears to be a secondary overprint (VRM), with the *in situ* direction oriented subparallel to the present field. This observation is also reflected in the NRM/IRM demagnetization results that show ratio values below 10⁻³ (Fig. 7). In contrast, representative specimens from LZ12 and 21 have NRM/IRM values that sit midway between a clear primary (10⁻²) and a clear secondary (10⁻³). Site LZ12 has, in tilt-corrected coordinates, a NNW declination and a negative inclination (–25.0°), whereas LZ21 has a SSE declination and a negative, shallow (–9.0°) inclination. Interestingly, the palaeomagnetic behaviour (e.g. the presence of two components of magnetization) and the mean ChRM direction of LZ21 are similar to those of the pillow basalts. This observation further supports the notion that the pillow basalts and the dyke are magnetically coeval (hence, also implying a primary magnetization for the dyke). Combining the ChRM direction of LZ21 with the individual mean ChRM directions from four pillow basalt sites gives a direction of: *in situ* Dec = 162.2°, Inc = –21.4°, $\alpha_{95} = 17.0^\circ$, $k = 21.1$; tilt-corrected Dec = 159.3°, Inc = –12.5°, $\alpha_{95} = 6.0^\circ$, $k = 162.5$. These data suggest that the rocks were formed at a subequatorial location (6.3°N ± 3.1°). Passage of the fold test is determined by getting the ratio of k after correction (k_a) to k before tectonic correction (k_b) (Butler 1992). The ratio obtained for the Chico River samples is 7.20. This k_a/k_b value exceeds the value of F distribution for the 5 per cent significance level ($F(6,6)$) of 4.28. The result suggests the statistical passage of the fold test.

As for LZ12, there is no compelling evidence to suggest that the site carries a primary remanence. In fact, the tilt-corrected mean direction of LZ12 resembles the LCC direction of some specimens from the pillow basalts and hence, implying that the magnetizations of LZ12 specimens are secondary overprints.




Figure 4. Summary of palaeomagnetic directional data from ophiolitic basement rocks along the Chico River and adjacent areas. Upper plots are the low coercivity component (LCC) directions of specimens ($MAD < 10^\circ$). Lower plots are the characteristic remanent magnetization (ChRM) mean directions shown with their 95 per cent confidence circles. Solid/open symbols are downward/upward directed.

IRM acquisition experiments followed by the thermal demagnetization of IRM of a specimen from LZ21 indicate magnetite and some maghemite as principal carriers of remanence, similar to those noted in pillow basalt specimens (Fig. 6).

6 DISCUSSION

The Chico River pillow basalts provide evidence for the existence of a Mesozoic–early Cenozoic oceanic substratum upon which Luzon and neighbouring regions within the PMB were likely built (Queaño *et al.* 2008). Other Mesozoic ophiolitic outcrops in southeastern Luzon, such as the Upper Jurassic–Lower Cretaceous Lagonoy Ophiolite (Geary *et al.* 1988; Billedo 1994; Tamayo *et al.* 1996; David *et al.* 1997) and the ? Upper Cretaceous Angat Ophiolite (Haeck 1987; Arcilla *et al.* 1989; Yumul 1993), provide additional evidence. Similar outcrops can also be found in the central Philippines (south of Luzon) that include Samar, Antique, Cebu, Bohol, Dinagat and eastern Mindanao (McCabe & Almasco 1985; Hashimoto *et al.* 1979; Geary *et al.* 1988; Rangin 1990; Louca 1992). Geochemical studies (e.g. Yumul *et al.* 1992; Billedo 1994; Tamayo *et al.* 2001; Faustino *et al.* 2006; Yumul *et al.* 2006) indicate a suprasubduction signature for the ophiolites.

In the other regions of the western Pacific, pieces of Mesozoic suprasubduction ophiolite are also scattered (Pubellier *et al.* 2003). Ballantyne (1991) and Ali *et al.* (2001) reported Mesozoic ophiolite

basement on eastern Halmahera, as well as on the islands of Gag, Gebe, Waigeo, Obi and other islands along the Sorong fault zone. Radiometric dates and fossil evidence indicate an age varying from Middle to Late Jurassic (Ballantyne 1991) to Early Cretaceous or older (Pigram & Panggabean 1984; Ali *et al.* 2001) for the ophiolites.

The similarity in both age and geochemistry of the ophiolite rocks in the western Pacific region suggests that these rocks might have a common provenance or basin origin. Unfortunately, much of the pre-Cenozoic tectonic elements in the region have been destroyed, thus, making the tectonic reconstruction for the region prior to the Cenozoic problematic. In the Philippines, obtaining palaeomagnetic data from Mesozoic ophiolitic rocks for tectonic reconstruction have met with only limited success. Previously acquired palaeomagnetic data from these units (e.g. Angat Ophiolite) are not only sparse but also unreliable owing to wide scatter of the palaeomagnetic plots (McCabe *et al.* 1987). The most reliable pre-Cenozoic palaeomagnetic data come from the terranes of Palawan, Mindoro and Panay in the Central Philippines (McCabe *et al.* 1987). However, these terranes are ‘exotic’ and of continental origin, having been derived from southern China (Taylor & Hayes 1983; McCabe *et al.* 1987; Yumul *et al.* 2003).

Nevertheless, despite the relatively few data obtained from this study, there are strong indications based on available regional geologic and other previous data regarding the conclusions presented here. Taking the palaeomagnetic data from previous authors (e.g. Queaño *et al.* 2007), showing Luzon essentially located at lower latitudes during most of the Cenozoic (at least starting from Eocene time), it is very likely that the Chico pillow basalts also originated near these latitudes, given the time frame involved in the formation of these basalts (i.e. Cretaceous–Eocene based on inferred age).

The primary magnetization of the Chico River pillow basalts and dykes provide clues on the Mesozoic–early Cenozoic evolution of northern Philippines. The data suggest that the ophiolitic basement of the Central Cordillera was formed at subequatorial latitudes ($6.3^\circ\text{N} \pm 3.1^\circ$). Determining the amount and direction of rotation the basement rocks had undergone since its formation, however, is difficult to determine given the complex Cenozoic deformation that affected Luzon. Interestingly, Ali *et al.* (2001) obtained palaeomagnetic data from two areas, where a Mesozoic ophiolite on Obi Island in eastern Indonesia is exposed. Their results also suggest formation at a subequatorial latitude (mean tilt-corrected direction for the two areas is $\text{Dec} = 219.4^\circ$, $\text{Inc} = 12.1^\circ$, but the angular separation of 20.1° is noted).

Tracing the plate affinity of the ophiolitic basement of the Central Cordillera and other regions in Luzon is still an enigma. Using palaeomagnetic data as principal evidence, Queaño *et al.* (2007) suggest that during the early Cenozoic, northern Luzon formed part of the Philippine Sea Plate and had also been closer to the equator. Given the results herein presented, this implies that the ophiolitic basement rock of the Central Cordillera originated not too far from where the northern Luzon and adjacent regions were during their early stage of evolution. Deducing from Queaño *et al.*'s (2007) reconstruction, it would also be likely that the ophiolite basement (including the Chico basalts) of Luzon could have been part of the Philippine Sea Plate. Note, however, that models regarding the origin of the Cretaceous ophiolites in the Philippines point to either an autochthonous (e.g. Encarnacion 2004; Tamayo *et al.* 2004; Yumul 2007) or an allochthonous nature for these complexes. The latter model suggests that Luzon's ophiolitic basement is composite, the units being amalgamated by various tectonic processes, including subduction-related collision and strike-slip faulting

Table 1. Summary of palaeomagnetic data from the ophiolitic basement rocks along the Chico River and adjacent areas in the Central Cordillera, northern Luzon, Philippines.

Sample No.	Lithology	NRM (nT A J m)	N_p/N_c	LCC						ChRM							
				In situ			Tilt-corrected			In situ			Tilt-corrected				
				Dec	Inc	α_{95}	Dec	Inc	α_{95}	Dec	Inc	α_{95}	Dec	Inc	α_{95}		
Locality—Chico-1: Lat = 17° 12.638'N, Long = 121° 01.723'E; Strike = 018°, Dip = 50° → SE																	
LZ11	Pillow basalts	Site rejected; results overprinted by LZ12															
LZ12	Diabase dyke	334–402															
Locality—Chico-2: Lat = 17° 07.457'N, Long = 121° 02.548'E; 228°, Dip = 45° → NW																	
LZ13	Diabase dyke	73–377															
LZ14	Pillow basalt	274–3679	3/6	328.6	24.7	328.1	-19.6	67.8	4.4	4/6	338.5	21.4	355.8	-25.4	11.2	68.5	0.99
Locality—Chico-3: Lat = 17° 09.249'N, Long = 121° 02.895'E; Strike = 168°, Dip = 34° → W																	
LZ15	Pillow basalt	0.1–3	3/4	352.3	12.7	344.2	13.7	17.2	52.5	5/6	167.4	-15.6	159.2	-13.3	6.9	125.0	0.99
LZ16	Pillow basalt	1–86	3/6	353.2	26.4	336.1	25.6	16.1	59.3	3/6	168.6	-20.8	156.3	-18.6	5.0	6027	0.99
Locality—Chico-4: Lat = 17° 09.639'N, Long = 121° 02.994'E; Strike = 135°, Dip = 15° → SW																	
LZ17	Pillow basalts	0.7–1.1	4/7	331.6	17.8	326.3	21.9	36.8	7.2	7/7	162.8	3.0	162.7	-4.4	3.6	275.7	0.99
LZ18	Diabase dyke	Site rejected; no common trend															
Locality—Chico-5: Lat = 17° 12.136', Long = 121° 04.204'; Strike = 240°, Dip = 20° → NW																	
LZ19	Pillow basalt	Site rejected; only two useful specimen															
LZ20	Pillow basalt	Site rejected; no common trend															
LZ21	Diabase dyke	3–54	6/7	5.3	13.6	5.2	-11.2	9.9	46.6	6/7	153.4	-32.4	156.8	-9.2	4.8	196.9	0.99
LZ22	Pillow basalt	0.7–1	3/6	356.3	162	3564	-8.9	150	20.9	3/6	156.7	-40.5	161.2	-16.8	9.8	159.8	0.99
ChRM Mean																	
Pillow basalts (U15, 16, 17 and 22)																	
Mean <i>in situ</i>																	
Mean tilt-corrected																	
Pillow basalts and dyke (LZ15, 16, 17, 21 and 22)																	
Mean <i>in situ</i>																	
Mean tilt-corrected																	




Figure 5. IRM acquisition curves for representative specimens from ophiolitic rock sites along the Chico River and adjacent areas.




Figure 6. Thermal demagnetization of a three-component isothermal remanent magnetization (IRM; Lowrie 1990) for representative specimens from ophiolitic rock sites along the Chico River and adjacent areas.

(e.g. Dimalanta & Yumul 2004). In addition, previous workers (e.g. Yumul *et al.* 2003; Tamayo *et al.* 2004; Dimalanta & Yumul 2006) further suggested that these ophiolites could have been part of the proto-Philippine Sea Plate or proto-Molucca Sea Plate of Cretaceous age. Whether these ‘proto plates’ did exist is still a subject

of debate. To note, Ali *et al.* (2001) implied that the Philippine Sea Plate was already existent since the Mesozoic.

Clearly, the various models relating to the origin of the Cretaceous ophiolites point to the limited information on these rocks. The palaeomagnetic data herein presented provide crucial




Figure 7. NRM versus IRM demagnetization curves for representative specimens from ophiolitic basement rocks along the Chico River and adjacent areas.

information for interpreting the nature and origin of these complexes. Unfortunately, this data set is very small to allow plate reconstructions and modelling (including that for northern Luzon) for this period.

7 CONCLUSIONS

The new data from the Chico River pillow basalts and dykes presented here suggest that ophiolite basement rocks of northern Luzon formed at subequatorial latitudes. Conversely, together with those previously gathered from the island, the data suggest that northern Luzon was close to the equator from as long ago as the Cretaceous (or even earlier) to early Cenozoic. Additional palaeomagnetic data are needed to decipher the basinal/plate affinity of these rocks and to allow reconstructions for the region during the Mesozoic. This situation can be remedied through palaeomagnetic studies on the Mesozoic rocks within the Philippine archipelago and surrounding regions. Potential targets include the Angat Ophiolite and the Lagonoy ophiolite in southeastern Luzon, as well as the Mesozoic ophiolite outcrops in the central and southern Philippines. Already, there is a large set of information available on the geology of these regions that can be used to support palaeomagnetic studies.

ACKNOWLEDGMENTS

This work and the postgraduate studentship of KLQ were supported by the University of Hong Kong. The Mines and Geosciences Bureau, Philippines, provided the logistical support. Reviews by E. Appel, M. Fuller and an anonymous reviewer helped improve the manuscript. Constructive insights were offered by D. Faustino-Eslava and the members of the Rushurgent Working Group, National Institute of Geological Sciences.

REFERENCES

- Ali, J.R., 1989. Magnetostratigraphy of early Paleogene sediments from NW Europe, *PhD thesis*. University of Southampton, Southampton, 277 pp.
- Ali, J.R. & Hall, R., 1995. Evolution of the boundary between the Philippine Sea Plate and Australia, *Tectonophysics*, **251**, 251–275.
- Ali, J.R., Hall, R. & Baker, S.J., 2001. Paleomagnetic data from a Mesozoic Philippine Sea Plate ophiolite on Obi Island, Eastern Indonesia, *J. Asian Earth Sci.*, **19**, 537–548.
- Arcilla, C.A., Ruelo, H.B. & Umbal, J., 1989. The Angat ophiolite, Luzon, Philippines: lithology, structure, and problems in age interpretation, *Tectonophysics*, **168**, 127–135.
- Aurelio, M.A., 2000. Shear partitioning in the Philippines: constraints from Philippine Fault and global positioning system data, *Island Arc*, **9**, 584–597.
- Ballantyne, P., 1991. Petrological constraints upon the provenance and genesis of the East Halmahera ophiolite, *J. SE Asian Earth Sci.*, **6**, 259–269.
- Barrier, E., Huchon, P. & Aurelio, M.A., 1991. Philippine Fault: a key to Philippine kinematics, *Geology*, **19**, 32–35.
- Bautista, C.B., Bautista, M.L.P., Oike, K., Wu, F.T. & Punongbayan, R.S., 2001. A new insight on the geometry of subducting slabs in northern Luzon, Philippines, *Tectonophysics*, **339**, 279–310.
- Bellon, H. & Yumul, G.P., Jr., 2000. Mio-Pliocene magmatism in the Baguio Mining District (Luzon, Philippines): age clues to its geodynamic setting, *C.R. Acad. Sci. Paris, Sci. de la Terre et des planets*, **331**, 295–302.
- Billedo, E.B., 1994. Geologie de la Sierra Madre Septentrionale et de l'Archipel de Polillo (Ceinture Mobile Est Pacifique), *PhD thesis*. Nice-Sophia Antipolis Universite, France, 215 pp.
- Butler, R.F., 1992. *Paleomagnetism: Magnetic Domains to Geologic Terranes*, Blackwell Scientific Publications, Boston, 319 pp.
- Cardwell, R.K., Isack, B.L. & Karig, D.E., 1980. The spatial distribution of earthquakes, focal mechanism solutions, and subducted lithosphere in the Philippine and northeastern Indonesian islands, in *The Tectonic and Geological Evolution of SE Asian Seas and Islands*, *Geophysical Monograph*, Vol. **23**, pp. 1–35, ed. Hayes, D.E., American Geophysical Union.
- Cisowski, S.M., Dunn, J.R., Fuller, M. & Wasilewski, P.J., 1990. NRM : IRM(s) demagnetization plots of intrusive rocks and the origin of their NRM, *Tectonophysics*, **184**, 35–54.
- David, S., Jr., Stephan, J.F., Delteil, J., Muller, C., Butterlin, J., Bellon, H. & Billedo, E., 1997. Geology and tectonic history of Southeastern Luzon, Philippines, *J. Asian Earth Sci.*, **15**, 435–452.
- Dimalanta, C.B. & Yumul, G.P., Jr., 2004. Crustal thickening in an active margin setting (Philippines): the whys and the hows, *Episodes*, **27**, 260–264.
- Dimalanta, C.B. & Yumul, G.P., Jr., 2006. Magmatic and amagmatic contributions to crustal growth in the Philippine island arc system: comparisons of the Cretaceous and post-Cretaceous periods, *Geosci. J.*, **10**, 321–329.
- Encarnacion, J., 2004. Multiple ophiolite generation preserved in the northern Philippines and the growth of an island arc complex, *Tectonophysics*, **292**, 103–130.
- Fisher, R.A., 1953. Dispersion on a sphere, *Proc. R. Soc. London*, **A217**, 295–305.
- Fitch, T.J., 1972. Plate convergence, transcurrent faults, and internal deformation adjacent to southeast Asia and the western Pacific, *J. geophys. Res.*, **77B**, 4432–4460.
- Florendo, F.F., 1994. Tertiary arc rifting in northern Luzon, Philippines, *Tectonics*, **13**, 623–640.
- Fuller, M., McCabe, R., Williams, S., Almasco, J., Encina, R.Y. & Zanolina, A.S., 1983. Paleomagnetism of Luzon, in *The Tectonic and Geological Evolution of SE Asian Seas and Islands*, *Geophysical Monograph*, Vol. **27**, pp. 79–94, ed. Hayes, D.E., American Geophysical Union.
- Fuller, M., Cisowski, S., Hart, M., Haston, R., Schmidtke, E. & Jarrard, R., 1988. NRM:IRM(s) demagnetization plots—an aid to the interpretation of natural remanent magnetization, *Geophys. Res. Lett.*, **15**, 518–521.
- Faustine, D.V., Yumul, G.P. Jr., Dimalanta, C.B., De Jesus, J.V., Zhou, M.-F., Alchison, J. & Tamayo, R.A. Jr., 2006. Volcanic-hypabyssal rock

- geochemistry of a subduction-related marginal basin ophiolite: south-east Bohol Ophiolite-Cansiwang Mélange Complex, Central Philippines, *Ceascl. J.*, **10**, 291–303.
- Geary, E.E., Harrison, T.M. & Heizler, M., 1988. Diverse ages and origins of basement complexes, Luzon, Philippines, *Geology*, **16**, 341–344.
- Haecck, G.D., 1987. The geological and tectonic history of the central portion of the southern Sierra Madre, Luzon, Philippines, *PhD thesis*. 294 pp., Cornell University.
- Halls, H.C., 1978. The use of converging remagnetization circles in palaeomagnetism, *Phys. Earth and Planet. Int.*, **16**, 1–111.
- Hashimoto, W.E., Kitamura, N., Balce, G.R., Matsumaru, K., Kurihara, K. & Aliate, E.Z., 1979. Large foraminifera from the Philippines, part X: stratigraphic and faunal breaks between the Maybangan and Kinabuan formations in the Tanay region, Rizal, Philippines, *Geol. Paleontol. SE Asia*, **20**, 143–158.
- International Geomagnetic Reference Field (IGRF), 2005. 10th Generation, <http://www.ngdc.noaa.gov/AGA/vmod/igrf.html>.
- Kirschvink, J.L., 1980. The least square line and plane and the analysis of paleomagnetic data, *Geophys. J. R. astr. Soc.*, **62**, 699–718.
- Lewis, S.D. & Hayes, D.E., 1984. A geophysical study of the Manila Trench, Luzon, Philippines, *J. geophys. Res.*, **89B**, 9196–9214.
- Louca, K., 1992. Geology and mineral resources of Dinagat island, Exploration Area 2, Internal Report 2, United Nations Revolving Fund for Natural Resources Exploration, Project PHI/86/1, 23 pp.
- Lowrie, W., 1990. Identification of ferromagnetic minerals in a rock by coercivity and unblocking temperature properties, *Geophys. Res. Lett.*, **17**, 159–162.
- Maletterre, P., 1989. Contribution à l'étude géologique de la plaque de la mer des Philippines. Histoire sédimentaire, magmatique et métallogénique d'un arc cénozoïque déformé en régime de transpression, *PhD thesis*. Université de Bretagne Occidentale, France, 367 pp.
- McCabe, R.J. & Almasco, J.N., 1985. Terranes in the Central Philippines, in *Tectonostratigraphic Terranes of the Circum-Pacific Region*, pp. 421–436, ed. Howell, D.G.
- McCabe, R.J., Kikawa, E., Cole, J.T., Malicse, A.J., Baldauf, P.E., Yumul, G.P., Jr. & Almasco, J., 1987. Paleomagnetic results from Luzon and the Central Philippines, *J. geophys. Res.*, **92B**, 555–580.
- Ozawa, A., Tagami, T., Listanco, E.L., Arpa, C.B. & Sudo, M., 2004. Initiation and propagation of subduction along the Philippine Trench: evidence from the temporal and spatial distribution of volcanoes, *J. Asian Earth Sci.*, **23**, 105–111.
- Pena, R., 1992. A review of the stratigraphy of Baguio district, *J. geol. Soc. Phil.*, **47**, 151–166.
- Pigram, C.J. & Panggabean, H., 1984. Rifting of the northern margin of the Australian continent and the origin of some microcontinents in eastern Indonesia, *Tectonophysics*, **107**, 331–353.
- Pubellier, M., Quebral, R., Rangin, C., Deffontaine, B., Muller, C. & Butterlin, J., 1991. The Mindanao collision zone: a soft collision event within a continuous Neogene strike-slip setting, *J. SE Asian Earth Sci.*, **6**, 239–248.
- Pubellier, M., Ali, J. & Monnier, C., 2003. Cenozoic Plate interaction of the Australia and Philippine Sea Plates: 'hit-and-run' tectonics, *Tectonophysics*, **363**, 181–199.
- Queaño, K.L., 2006. Tectonic modeling of Northern Luzon, Philippines and regional implications, *PhD thesis*. University of Hong Kong, 498pp.
- Queaño, K.L., Ali, J.R., Milsom, J., Aitchison, J.C. & Pubellier, M., 2007. North Luzon and the Philippine Sea Plate motion model: insights following paleomagnetic, structural, and age-dating investigations, *J. geophys. Res.*, **112**, 1–44.
- Queaño, K.L., Ali, J.R., Aitchison, J., Yumul, G.P., Jr., Pubellier, M. & Dimalanta, C.B., 2008. Geochemistry of Cretaceous to Eocene ophiolitic rocks of the Central Cordillera: implications on the Mesozoic-early Cenozoic evolution of northern Philippines, *Int. geol. Rev.*, **50**, 407–421.
- Quebral, R., Pubellier, M. & Rangin, C., 1996. The onset of movement on the Philippine fault in eastern Mindanao: a transition from a collision to a strike slip environment, *Tectonics*, **15**, 713–726.
- Rangin, C., 1989. The Sulu Sea, a back-arc setting within a Neogene collision zone, *Tectonophysics*, **161**, 119–141.
- Rangin, C., 1990. The Philippine Mobile Belt: a complex plate boundary, *J. SE Asian Earth Sci.*, **6**, 209–220.
- Rangin, C. & Pubellier, M., 1990. Subduction and accretion of Philippine Sea Plate fragments along the Eurasian margin, in *Tectonics of Circum-Pacific Continental Margins*, pp. 139–164, eds Auboin, J., Bourgois, J., Brill Academic Publishers, Inc., Leden, The Netherlands.
- Ringenbach, J.C., 1992. La Faille Philippine et les chaînes en décrochement associées (center et nord de Luzon): evolution Cenozoïque et cinématique des déformations quaternaires, *PhD thesis*. Université de Nice Sophia Antipolis, France, 336 pp.
- Ringenbach, J.C., Pinet, N., Stephan, J.F. & Delteil, J., 1993. Structural variety and tectonic evolution of strike-slip basins related to the Philippine Fault System, Northern Luzon, Philippines, *Tectonics*, **12**, 187–203.
- Stephan, J.F., Blanchet, R., Rangin, C., Pelletier, B., Letouzey, J. & Muller, C., 1986. Geodynamic evolution of the Taiwan-Luzon-Mindoro belt since the late Eocene, *Tectonophysics*, **125**, 245–268.
- Tamayo, R.A., Jr., Yumul, G.P., Jr. & Jumawan, F.T., 1996. Geology and geochemistry of the mafic rock series of the Camarines Norte Ophiolite Complex, *J. geol. Soc. Phil.*, **51**, 131–152.
- Tamayo, R.A., Jr., Yumul, G.P., Jr., Maury, R.C., Polve, M., Cotten, J. & Bohn, M., 2001. Petrochemical investigation of the Antique Ophiolite (Philippines): implications on volcanogenic massive sulfide and podiform chromitite deposits, *Res. Geol.*, **51**, 145–164.
- Tamayo, R.A., Jr., Maury, R.C., Yumul, G.P., Jr., Polvé, M., Cotten, J., Dimalanta, C.B. & Olaguera, F., 2004. Subduction-related magmatic imprint of most Philippine ophiolites: implications on the early geodynamic evolution of the Philippine archipelago, *Bull. Soc. Geol. Fr.*, **175**, 443–460.
- Taylor, B. & Hayes, D.E., 1983. Origin and history of the South China Sea basin, in *The Tectonic and Geological Evolution of SE Asian Seas and Islands*, *Geophysical Monograph*, **Vol. 27**, pp. 23–56, ed. Hayes, D.E., American Geophysical Union.
- United Nations Development Programme (UNDP), 1987. Geology and mineralization in the Baguio area, Northern Luzon, UNDP Technical Report No. 5, pp. 1–82.
- Yumul, G.P., Jr., 1993. Angat Ophiolitic Complex, Luzon, Philippines: a Cretaceous dismembered marginal basin ophiolitic complex, *J. SE Asian Earth Sci.*, **8**, 529–537.
- Yumul, G.P., Jr., 1994. A marginal basin crust basement for the Baguio mining district, Luzon, Philippines, *J. geol. Soc. Phil.*, **49**, 79–87.
- Yumul, G.P., Jr., 2007. Westward younging disposition of Philippine ophiolites and its implications for arc evolution, *Island Arc*, **16**, 306–317.
- Yumul, G.P., Jr., Samson, M.T., Claveria, R.J.T., de Silva, L.P., Jr., Diomampo, E.O. & Rafols, J.P., 1992. Preliminary geochemical evidences for a marginal basin basement complex for the Baguio Mining District, Luzon, Philippines, *J. geol. Soc. Phil.*, **47**, 5–17.
- Yumul, G.P., Jr., Dimalanta, C.B., Tamayo, R. & Maury, R.C., 2003. Collision, subduction and accretion events in the Philippines: a synthesis, *Island Arc*, **12**, 77–91.
- Yumul, G.P., Jr., Dimalanta, C.B., Tamayo, R.A., Jr. & Zhou, M.F., 2006. Geology and geochemistry of the Rapu-Rapu ophiolite complex, eastern Philippines: possible fragment of the proto-Philippine Sea Plate, *Int. geol. Rev.*, **48**, 329–348.
- Yumul, G.P., Jr., Dimalanta, C.B., Marquez, E.J. & Maglambayan, V.B., 2008. Tectonic setting of a composite terrane: a review of the Philippine island arc system, *Geosci. J.*, **12**, 7–17.
- Zijderveld, J.D.A., 1967. A.C. demagnetization of rocks, in *Methods in Paleomagnetism*, pp. 256–286, eds Collinson, D.W., Creer, K.M. & Runcorn, S.K., Elsevier, New York.