BIOCHEMICAL CHARACTERISTICS OF A DIFFUSIBLE FACTOR THAT INDUCES GAMETOPHYTE TO SPOROPHYTE SWITCHING IN THE BROWN ALGA ECTOCARPUS

Haiqin Yao, Delphine Scornet, Murielle Jam, Cécile Hervé, Philippe Potin,
Lydie Oliveira Correia, Susana M Coelho, J. Mark Cock

► To cite this version:

Haiqin Yao, Delphine Scornet, Murielle Jam, Cécile Hervé, Philippe Potin, et al.. BIOCHEMICAL CHARACTERISTICS OF A DIFFUSIBLE FACTOR THAT INDUCES GAMETOPHYTE TO SPOROPHYTE SWITCHING IN THE BROWN ALGA ECTOCARPUS. *Journal of Phycology*, 2021, 57 (3), pp.742-753. 10.1111/jpy.13126 . hal-03089431

HAL Id: hal-03089431

<https://cnrs.hal.science/hal-03089431>

Submitted on 28 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**BIOCHEMICAL CHARACTERISTICS OF A DIFFUSIBLE FACTOR THAT
INDUCES GAMETOPHYTE TO SPOROPHYTE SWITCHING IN THE
BROWN ALGA *ECTOCARPUS***

*Haiqin Yao¹, Delphine Scornet¹, Murielle Jam², Cécile Hervé², Philippe Potin³, Lydie Oliveira
Correia⁴, Susana M. Coelho^{1,*}, J. Mark Cock^{1,*}*

¹Algal Genetics Group, CNRS, Sorbonne Université, UPMC University Paris 06, UMR 8227, Integrative
Biology of Marine Models, Station Biologique de Roscoff, CS 90074, F-29688, Roscoff, France

²Marine Glycobiology, CNRS, Sorbonne Université, UPMC University Paris 06, UMR 8227, Integrative
Biology of Marine Models, Station Biologique de Roscoff, CS 90074, F-29688, Roscoff, France

³Algal Biology and Environmental Interactions, CNRS, Sorbonne Université, UPMC University Paris 06,
UMR 8227, Integrative Biology of Marine Models, Station Biologique de Roscoff, CS 90074, F-29688,
Roscoff, France

⁴Micalis Institute, PAPPSO, INRA, AgroParisTech, Université Paris-Saclay, 78350, Jouy-en-Josas, France

* Authors for correspondence (cock@sb-roscoff.fr, coelho@sb-roscoff.fr, tel.: 33 (0)2 98 26 23 60), joint last
authors.

Running title: *Ectocarpus* sporophyte-inducing diffusible factor

ABSTRACT

The haploid-diploid life cycle of the filamentous brown alga *Ectocarpus* involves alternation between two independent and morphologically distinct multicellular generations, the sporophyte and the gametophyte. Deployment of the sporophyte developmental program requires two TALE homeodomain transcription factors OUROBOROS and SAMSARA. In addition, the sporophyte generation has been shown to secrete a diffusible factor that can induce uni-spores to switch from the gametophyte to the sporophyte developmental program. Here, we determine optimal conditions for production, storage and detection of this diffusible factor and show that it is a heat-resistant, high molecular weight molecule. Based on a combined approach involving proteomic analysis of sporophyte-conditioned medium and the use of biochemical tools to characterize arabinogalactan proteins, we present evidence that sporophyte-conditioned medium contains AGP epitopes and suggest that the diffusible factor may belong to this family of glycoproteins.

Key index words: arabinogalactan protein, diffusible factor, *Ectocarpus*, gametophyte, life cycle, sporophyte

Abbreviations: ACN, acetonitrile; AGP, arabinogalactan protein; LC-MS/MS, liquid chromatography coupled to mass spectrometry; ORO, OUROBOROS; PES, Provasoli-enriched natural seawater; SAM, SAMSARA; SCM, sporophyte-conditioned medium; SDS-PAGE, sodium dodecyl sulfate–polyacrylamide gel electrophoresis; TALE HD TF, three amino acid loop extension homeodomain transcription factor; TFA, trifluoroacetic acid, WSC, wall sensing component.

INTRODUCTION

Most eukaryotic life cycles involve an alternation between haploid and diploid phases, with the transitions between ploidy states occurring as a result of meiotic division (ploidy reduction) and gamete fusion, or syngamy (increased ploidy; Coelho et al. 2007). Multicellular photosynthetic organisms with haploid-diploid life cycles produce two distinct multicellular generations, one during each phase of the life cycle: diploid, spore-producing sporophytes and haploid, gamete-producing gametophytes (Coelho et al. 2007, Cock et al. 2014). The majority of brown algae have this type of haploid-diploid life cycle and one of these species, the filamentous brown alga *Ectocarpus* sp., is being used as a model system to study life cycle regulation (Peters et al. 2004, Cock et al. 2014, Coelho and Cock 2020). *Ectocarpus* has a complex life cycle (Müller 1967). In addition to the basic sexual life cycle, involving an alternation between gametophyte and sporophyte

56 generations, several asexual variations have been observed in culture, including parthenogenetic
 57 development of gametes that fail to fuse with a gamete of the opposite sex (Müller 1967). Interestingly,
 58 germinating parthenogenetic gametes deploy the sporophyte program, despite being haploid, to produce
 59 partheno-sporophyte individuals that are morphologically indistinguishable from diploid sporophytes.
 60 Moreover, these individuals not only resemble sporophytes morphologically but are functional sporophyte
 61 thalli, producing spores and not gametes (Müller 1967, Bothwell et al. 2010). One important conclusion that
 62 can be drawn from the existence of haploid partheno-sporophytes is that life cycle generation (i.e.
 63 deployment of a gametophyte or a sporophyte bodyplan) is not determined by ploidy (i.e. whether the
 64 organism is in the haploid or diploid phase) and that these two features of the life cycle can be uncoupled
 65 under certain circumstances (Müller 1967, Bothwell et al. 2010). This conclusion is supported by the
 66 existence of genetic mutants that cause switching between life cycle generations, independent of the ploidy
 67 of the mutant individual (Coelho et al. 2011, Arun et al. 2019). Genetic analysis of life cycle mutants of this
 68 type has demonstrated that the deployment of the sporophyte program in *Ectocarpus* is under the control of
 69 two homeodomain transcription factors (HD TFs) of the three amino acid loop extension (TALE) class,
 70 which have been named OUROBOROS (ORO) and SAMSARA (SAM), respectively (Coelho et al. 2011,
 71 Arun et al. 2019). These transcription factors appear to be derived from an extremely ancient life cycle
 72 regulation system and to be distantly related to HD TF life cycle regulators in other eukaryotic supergroups
 73 such as specific KNOX and BEL class proteins in the green lineage (Viridiplantae).

74 Further evidence for the independence of life cycle generation and ploidy was provided by the identification
 75 of a sporophyte-inducing factor that is secreted into the surrounding seawater medium by *Ectocarpus*
 76 sporophytes in culture (Arun et al. 2013). When uni-spores (i.e. spores produced by unilocular sporangia),
 77 which normally develop to produce the gametophyte generation, are allowed to germinate in the presence of
 78 this diffusible factor, a proportion of the resulting germlings deploy the sporophyte developmental pathway.
 79 Interestingly, uni-spores carrying either *oro* or *sam* mutations are resistant to the action of the diffusible,
 80 sporophyte-inducing factor (Arun et al. 2013, 2019), indicating that the *ORO* and *SAM* genes are necessary
 81 for developmental reprogramming to occur. *ORO* and *SAM* may therefore be part of the regulatory network
 82 triggered by the sporophyte-inducing factor.

83 The objective of this study was to further characterise the diffusible sporophyte-inducing factor. After
 84 optimising production, storage and bioassay of the factor, we carried out a number of analyses aimed at
 85 providing further information about its biochemical nature. We obtained evidence that the sporophyte-
 86 inducing factor is a high molecular weight molecule that is resistant to high temperature and protease
 87 treatment. Interestingly, mass spectrometry analysis of partially purified (ultrafiltrated) sporophyte-
 88 conditioned medium (SCM) identified 36 proteins, one of which is predicted to contain an arabinogalactan
 89 protein (AGP) core protein domain. Furthermore, AGP glycan epitopes were detected in a concentrated SCM
 90 preparation using immunoblotting and SCM activity was reduced following incubation with an AGP-reactive

Yariv reagent. Taken together, these observations led us to suggest that the diffusible factor may correspond to an AGP.

MATERIALS AND METHODS

Biological material and preparation of SCM. The biological material used in this study corresponded to partheno-sporophytes and gametophytes of *Ectocarpus* sp. strain Ec32, which is a male descendant of a diploid sporophyte Ec17 (CCAP1310/193) isolated at San Juan de Marcona, Peru (Peters et al. 2008). The original Ec32 gametophyte was derived from a uni-spore produced in a unilocular sporangium on the diploid sporophyte parent strain Ec17. In the unilocular sporangia of diploid sporophytes, uni-spores are produced by a single meiotic division followed by several mitotic divisions. The spores can therefore be termed uni-spores (derived from a unilocular sporangium) or meio-spores (derived meiotically). Parthenogenetic development of male gametes, produced in plurilocular gametangia on the original Ec32 gametophyte, gave rise to the Ec32 partheno-sporophyte strain, which is maintained in culture by asexual reproduction either by thallus fragmentation or through mitotically-derived mito-spores produced in plurilocular sporangia. Ec32 gametophytes were derived from Ec32 partheno-sporophytes through the production of uni-spores in unilocular sporangia. A previous study (Bothwell et al. 2010) showed that partheno-sporophytes can produce uni-spores in two different ways: either the partheno-sporophyte undergoes endoreduplication during early development to become diploid, allowing normal meiotic divisions to occur in the unilocular sporangia or the thallus remains haploid, in which case the first division of the uni-spore mother cell is apomeiotic (a meiotic division without reduction). Both of these processes produce haploid uni-spores which develop as gametophytes.

Strain Ec32 was cultivated in Provasoli-enriched natural seawater (PES) under standard culture conditions at 13°C, with a 12h/12h day/night cycle of white fluorescent light at 20 $\mu\text{M photons/m}^2\cdot\text{s}$ (Coelho et al. 2012). Strain Ec32 carries a V sex chromosome and is therefore male when in the gametophyte generation.

The standard conditions for the production of SCM (Yao et al. 2020) involved inoculating 160 ml of PES in a 150 mm Petri dish with about 10^6 gametes (which germinate to produce partheno-sporophytes) and cultivating for 4 to 6 weeks under low light conditions (2-3 $\mu\text{M photons/m}^2\cdot\text{s}$). Variations on this procedure were used to test the effect of light intensity, culture density and culture time on SCM production. The alternative light conditions used for these tests were medium light (15-20 $\mu\text{M photons/m}^2\cdot\text{s}$) and high light (25-30 $\mu\text{M photons/m}^2\cdot\text{s}$).

To produce large volumes of SCM, 10 L bottles of PES were inoculated with 0.5 g of partheno-sporophyte material and cultured under low light conditions with aeration by pumping air through a 0.2 μm filter. In this bottle system, cultures grown for between four and 11 weeks produced SCM with diffusible factor activity.

SCM was recovered by filtering cultures through a reusable coffee filter (12.5 cm Finlandek permanent coffee filter) to remove most of the partheno-sporophyte tissue and then filtering through a 40 µm cell strainer (Falcon 40 µm Nylon cell strainer) to remove any remaining algal cells. Finally, a 0.22 µm syringe filter (Millipore Millex-GP polyethersulphone membrane) was used to remove bacteria. The SCM was then either used immediately or stored at 4°C.

Bioassay for the diffusible sporophyte-inducing factor. Test samples of SCM were either diluted in PES or, when used undiluted, enriched in micronutrients by addition of Provasoli solution to 1x final concentration. One or more unilocular sporangia were micro-dissected from fertile partheno-sporophyte thalli under a binocular microscope and placed in 300 µl of these preparations so that the uni-spores were released directly into the test solution. Each 300 µl droplet represented a replicate assay of the test solution. The preparations were incubated overnight in very high light (35 µmol photons /m²·s) to promote uni-spore release. Empty unilocular sporangia were removed after release of the uni-spores. The numbers of gametophytes and sporophytes were scored four to seven days later based on germling morphology (Peters et al. 2008) using an inverted light microscope (Olympus CKX41). In some cases when the uni-spores grew poorly, an additional 300 µl of the test preparation was added to the drop and culture continued.

Proteinase K and heat treatment of SCM. Heat-treated SCM was autoclaved at 121°C for 30 min. Proteinase-treated SCM was incubated with 100 µg/ml proteinase K at 37°C for 1 h and then autoclaved at 121°C for 30 min. Provasoli solution was added to the samples from both experiments to 1x final concentration before carrying out bioassays for the diffusible factor.

Size fractionation by ultrafiltration. SCM was pre-filtered through a 0.22 µm filtration system (Stericup and Steritop, 500 ml Millipore Express PLUSE 0.22 µm PES) before ultrafiltration. Ultrafiltration was then carried out using an ÄKTATM flux (GE Healthcare BioSciences Uppsala, Sweden) equipped with a 100 kDa, a 50 kDa or a 10 kDa cartridge. Batches of pre-filtered SCM, were gradually fed into the starting reservoir in order to maintain a constant volume of about 300 ml, which was cycled through the filter resin, with the filtrate being collected separately during the cycling. The 10-50 kDa size range was obtained by filtering the flow-through from a 50 kDa filter filtration through a 10 kDa filter as a second step, retaining the retentate. Concentrated retentates were designated uf-SCM for ultra-filtrated SCM. To assay the activity of the diffusible factor, uf-SCM preparations were diluted in PES to obtain a concentration equivalent to 1X SCM.

Immunodot assays with anti-AGP antibodies. Serial dilutions of each test sample (five five-fold dilution steps) were carried out and 1 µL aliquots of each dilution spotted onto nitrocellulose membranes

158 (Amersham) along with equivalent dilutions of a gum arabic solution (starting concentration 1 mg/ml) as a
159 positive control. The prints were allowed to dry and blocked with 5% of milk powder in phosphate buffered
160 saline solution (PBS/MP). The samples were then probed with the rat anti-AGP monoclonal antibodies (Plant
161 Probes, Leeds, UK) JIM8, JIM13, JIM16, JIM4, LM2, LM14 and MAC207 diluted 10-fold in PBS/MP
162 (Moller et al., 2007). After washing, anti-rat-HRP (diluted 1000-fold) was added in PBS/MP. After an
163 additional wash, antibody-epitope interactions were detected using a luminescent ECL substrate (BioRad)
164 and a Fusion FX.XT-820.EPI/20M system (Vilber Lourmat, France).

165

166 *Yariv reagent tests.* β -D-glucosyl (β Glc) or β -D-mannosyl (β Man) Yariv reagents (Biosupplies Pty Ltd,
167 Melbourne, Australia) were added to Provasoli-enriched SCM at final concentrations of 2 μ g/ml before
168 carrying out diffusible factor bioassays.

169

170 *SCM analysis by SDS-PAGE.* An initial volume of 200 ml SCM was concentrated using two Amicon®
171 Ultra-15 30 kDa centrifugal filters by sequentially centrifuging 15 ml aliquots for 8 min at 3500 rpm. Each of
172 the retentates (1.5 ml for each of the two filters) was then washed with 100 ml MilliQ H₂O by adding batches
173 of 15 ml and centrifuging each time for 8 min at 3500 rpm. The final retentate, a total of 2 ml from the two
174 filters was freeze-dried and redissolved in 10 μ l MilliQ H₂O and 2 μ l of 6x protein loading buffer (360 mM
175 Tris-HCl pH 6.8, 45% Glycerol, 9% SDS, 9% DTT, 0.12% Bromophenol Blue). Twelve microlitres of this
176 sample was heated to 100°C for 10 min before separation by sodium dodecyl sulphate polyacrylamide gel
177 electrophoresis (SDS-PAGE, 12% w/v; Brunelle and Green 2014) at 120 V. Gels were stained with
178 Coomassie blue for 1h then washed three times in 20% ethanol/10% acetic acid.

179

180 *Protein digestion and mass spectrometry analysis.* Protein bands were excised from an SDS-PAGE gel for
181 mass spectrometry analysis, which was carried out at the Paris Sud Ouest PAPPSO proteomics core facility
182 (<http://papso.inra.fr>). Gel pieces were washed three times with 50 μ l of 10% formic acid in 40% ethanol, and
183 50 μ l of 50 mM sodium bicarbonate (NH₄HCO₃) in 50% acetonitrile (ACN). The gel pieces were incubated
184 in 10 mM dithiothreitol at 56°C to reduce any disulphide bridges and the cysteines were alkylated with 50
185 mM iodoacetamide. In-gel digestion was carried out overnight at 37°C with 100 ng of trypsin in 50 mM
186 sodium bicarbonate. Peptides were extracted with 0.1% trifluoroacetic acid (TFA) in 40% ACN, dried in a
187 speed vac and redissolved in 20 μ l of 0.1% TFA in 2% ACN. Four microlitres of this preparation were
188 injected into the LC-MS/MS system.

189 LC-MS/MS analyses were carried out using an Ultimate 3000 RSLC system (Thermo Scientific) coupled to
190 a LTQ-orbitrap discovery mass spectrometer (Thermo Scientific) by a nanoelectrospray ion source. The
191 protein digest was injected and preconcentrated on a precolumn (Acclaim PepMap C18 particle 5 μ m size, 5

mm length, 300 μm i.d., Thermo Fisher Scientific) at 20 $\mu\text{l}/\text{min}$ with 0.08% TFA in 2% ACN for 2 min, followed by a separation on a reverse phase separating column (Acclaim PepMap RSLC nanoViper, C18 particle 2 μm size, 150 mm length, 75 μm i.d., Thermo Fisher Scientific). Buffers were 0.1% formic acid in 98% water (solvent A) and 0.1% formic acid in 80% ACN (solvent B). The peptides were eluted with a multi-step gradient from 1% to 35% of solvent B for 34.5 min at 300 nl/min for a total run of 42 min. MS scans were acquired in a mass range of 300-1400 m/z at a resolution of 15000 in the orbitrap analyser. The eight most intense ions were selected for CID MS/MS with a normalised collision energy of 35 in the ion trap.

All MS/MS spectra were searched against a database of deduced proteins based on version 2 of the *Ectocarpus* genome annotation (<https://bioinformatics.psb.ugent.be/orcae/overview/EctsiV2>) using the X!TandemPipeline (version 3.4.3), an open search engine developed by PAPPSO (<http://pappso.inra.fr/bioinfo/xtandempipeline/>). Precursor mass tolerance was 10 ppm and fragment mass tolerance was 0.5 Da. Data was filtered using the following criteria: a peptide E-value of <0.01 , a protein E-value of $<10\text{e-}4$ and a minimum of two identified peptides per protein.

RESULTS

Rapidly released uni-spores are more responsive to the sporophyte-inducing diffusible factor. Arun et al. (2013) demonstrated that *Ectocarpus* sporophytes secrete a diffusible factor that induces uni-spores to switch from the gametophyte to the sporophyte developmental program. With the objective of further characterising this diffusible factor, we first aimed to determine optimal conditions for its production and assay. The physiological state of an sporophyte can influence unilocular sporangium production and function, leading to developmental abnormalities of the sporangium and often delayed or dysfunctional release of uni-spores. To assess the effect of uni-spore quality on detection of the diffusible factor, we compared batches of unilocular sporangia that had released after 12-24 h, 24-48 h or >48 h of incubation in SCM to determine whether the time taken to release the spores influenced the proportion of uni-spores that were converted to the sporophyte generation (Fig. 1A). This analysis indicated that rapidly released uni-spores (i.e. within 24 h) were more sensitive to the diffusible factor than uni-spores whose release had been delayed (i.e. taking >48 h; Wilcoxon-Mann-Whitney test, $W = 1614.5$, $p\text{-value} = 0.0313$). Based on this observation, whenever possible, subsequent experiments were carried out with uni-spores that had been released within 48 h of transfer of the micro-dissected unilocular sporangia to the test conditions.

The sporophyte-inducing diffusible factor is stable when stored at 4°C or at -20°C. To test the stability of the sporophyte-inducing factor, SCM was stored at either 4°C or at -20°C and its activity tested after different time periods (Fig. 1B). The results of this analysis indicated that the factor was stable for at least eight weeks

226 at 4°C. Similarly, storage at -20° for one to five weeks did not result in a decrease in the activity of the
227 diffusible factor (Fig. 1C). We noted that SCM activity increased after storage for more than five weeks at
228 4°C but no increase in activity was not observed when the medium was stored at -20°C for the same period
229 of time.

230

231 *Effect of sporophyte culture conditions on production of the sporophyte-inducing diffusible factor.* We next
232 tested whether the conditions under which the sporophyte material was cultivated influenced production of
233 the diffusible factor. To test whether diffusible factor production was influenced by the length of time the
234 sporophyte material was maintained in culture, Provasoli-enriched natural seawater (PES) was inoculated
235 with gametes and the resulting partheno-sporophytes were cultivated for different times under standard
236 conditions. The SCM samples were removed after four, six, eight and ten weeks and assayed for diffusible
237 factor activity (Fig. 1D). This experiment indicated that SCM contained the highest activities after six to
238 eight weeks of culture, with the activity diminishing in older cultures.

239 To test the effect of light intensity, PES was inoculated with gametes and the resulting partheno-sporophytes
240 grown either under low (2-3 $\mu\text{mol photons /m}^2\cdot\text{s}$), medium (15-16 $\mu\text{mol photons /m}^2\cdot\text{s}$) or high light (25-30
241 $\mu\text{mol photons /m}^2\cdot\text{s}$) conditions. SCM was collected after four weeks of culture and assayed for the diffusible
242 factor (Fig. 1E). There were no significant differences between the detected levels of diffusible factor
243 produced by the cultures that were grown under the three light conditions (Wilcoxon-Mann-Whitney test, W
244 = 1461, 785.5 and 571.5, p -values = 0.6847, 0.0740 and 0.2480 for low light versus normal light, low light
245 versus high light and normal light versus high light, respectively).

246 Based on these experiments, cultivation times of five to six weeks under low light conditions (to limit the
247 growth rate of the cultures) were used to produce SCM for subsequent experiments.

248

249 *The diffusible factor is a large molecule.* As a first step towards investigating the biochemical nature of the
250 diffusible factor, we used ultrafiltration to estimate its molecular size. SCM was filtered through an ÄKTA™
251 flux ultrafiltration system using filters with different molecular size cut-offs to prepare different size
252 fractions (Fig. 2A). Bioassay testing of the different size fractions indicated that the diffusible factor had a
253 molecular size of greater than 50 kDa and that at least part of the activity had a molecular size of less than
254 100 kDa (Fig. 2B).

255

256 *Mass spectrometry analysis of proteins in concentrated SCM.* To further characterise the high molecular
257 weight fraction, 200 ml of SCM was processed with a Falcon filter system to concentrate >30 kDa
258 components. Coomassie blue staining of the concentrated SCM sample after separation on SDS-PAGE (Fig.
259 3A) revealed several high molecular weight bands of greater than 50 kDa (i.e. in the active size range

determined by ultrafiltration experiments, Fig. 2B). Liquid chromatography coupled to mass spectrometry (LC-MS/MS) analysis of the SDS-PAGE bands identified peptides corresponding to 36 different proteins (Table 1). Most of the proteins detected were predicted to be secreted proteins based on the presence of a putative signal peptide and several of the proteins could potentially play a role in cell to cell signalling. For example AGPs have important signalling roles in land plants and the list included a mannuronan C5-epimerase (locusID Ec-20_004700, Fig. 3B) that had been reported to contain three chimeric AGP backbone motifs (Hervé et al. 2016).

267

The sporophyte-inducing diffusible factor is resistant to autoclaving and proteinase K treatment. To characterise the diffusible factor further, we determined whether it was stable when treated with high temperature or subjected to protease digestion. SCM was either autoclaved or digested by incubation at 37°C in the presence of 100 µg/ml proteinase K followed by autoclaving to denature the enzyme. Fig. 4 shows that the diffusible factor was resistant to both autoclaving and proteinase K treatment. Surprisingly, the activity of the factor significantly increased following proteinase K treatment (Wilcoxon-Mann-Whitney test, $W = 45.5$, p -value = 0.03785).

This result of this experiment suggested that the diffusible factor may not be a protein but note that alternative interpretations of these results are possible. For example, the factor could be a complex molecule such as a glycoprotein, with the protein part of the molecule being either inaccessible to proteinase K or unnecessary for the factor's activity. This interpretation would be consistent with the diffusible factor corresponding to an AGP for example (based on the detection of Ec-20_004700 in the SCM). We therefore carried out some additional experiments aimed at further exploring the possibility that the SCM contains active AGP-like molecules.

282

Detection of AGP glycan epitopes in sporophyte-conditioned medium. Anti-AGP antibodies were used to assay for the presence of AGP-like proteins in SCM. Fig. 5 shows that AGP epitopes were detected in the high molecular weight SCM fraction after it had been concentrated by ultrafiltration. The strongest signals were obtained with the JIM13, JIM4 and MAC207 antibodies, but JIM16, LM14 and JIM8 also weakly detected epitopes. This experiment, therefore, confirmed that sporophytes secrete AGP-like molecules into the culture medium.

289

Decreased bio-activity of SCM in the presence of an AGP-reactive Yariv reagent. To determine whether AGP-like molecules have a role in the sporophyte-inducing activity of SCM, Yariv reagents were added to sporophyte-conditioned medium before testing for bioactivity. β -D-glucosyl and β -D-galactosyl Yariv reagents have been shown to react specifically with plant AGPs (Gane et al. 1995, Paulsen et al. 2014) and

inhibit their functions (Johnson et al. 2003, Tan et al. 2012, Hervé et al. 2016). Fig. 6 shows that addition of the β Glc Yariv reagent to SCM resulted in a significant decrease in activity (Wilcoxon-Mann-Whitney test, $W = 115$, p -value = 0.0369) in the uni-spore bioassay, whereas addition of an inactive analogue, β Man Yariv reagent, as a control did not. The results of this experiment were consistent with the diffusible factor being an AGP.

DISCUSSION

Genetic experiments have shown that the switch from the gametophyte to the sporophyte generation in *Ectocarpus* is controlled by two TALE HD TFs, OUROBOROS and SAMSARA (Coelho et al. 2011, Arun et al. 2019), but this switch can also be influenced by a non-cell autonomous, sporophyte-inducing factor that is secreted into the medium by the sporophyte generation (Arun et al. 2013). The experiments carried out here were aimed at further characterising this diffusible factor. We show that light intensity does not significantly influence production of the diffusible factor but at lower light intensities the algae grow more slowly allowing them to be maintained at higher densities for longer periods of time. *Ectocarpus* partheno-sporophytes normally become sexually mature after three to four weeks under normal light and, therefore, cultures cannot be maintained for long periods under these conditions. We also showed that SCM did not lose its activity when stored for 8 weeks at either 4°C or -20°C.

Further characterisation of the SCM indicated that the sporophyte-inducing factor was resistant to heat (121°C for 30 min) and proteinase K treatment and that the factor was a large molecular weight molecule, greater than 50 kDa in size. Mass spectrometry analysis identified 36 different proteins in concentrated SCM, including a mannuronan C5-epimerase (locusID Ec-20_004700) that has been reported to contain chimeric AGP backbone motifs (Hervé et al. 2016). Brown algal cell walls play an important role in conferring resistance to mechanical stress and protection from predators (Popper et al. 2011). They are mainly composed of alginates and sulphated fucans and therefore differ markedly in their composition from the cell walls of land plants (Hervé et al. 2016). Mannuronan C5-epimerase acts on alginate, altering its mechanical properties by converting β -D-mannuronate to its epimer β -L-guluronate (Nyvall et al. 2003, Fischl et al. 2016). The mannuronan C5-epimerase protein Ec-20_004700 is predicted to contain wall sensing component (WSC) domains, in addition to its catalytic domain. WSC domains have been associated with cell wall sensing (Ohsawa et al. 2017, Oide et al. 2019) and resistance to various stress conditions in fungi (Lodder et al. 1999, Lommel et al. 2004, Dupres et al. 2009). Recent studies indicate that fungal WSC proteins might contribute to protein anchoring and colonization of plant hosts through their ability to bind a variety of cell wall polysaccharides from plants and fungi including β -1,3-glucans, xylans and chitin (Oide et al. 2019, Wawra et al. 2019). The WSC domain family is one of the largest protein domain families in brown algae, with 444 members in *Ectocarpus* sp. strain Ec32 (Dittami et al. 2020). Three of the 28 mannuronan C5-epimerase enzymes in *Ectocarpus* are predicted to possess WSC domains and, while their biochemical

function has not been yet elucidated, these motifs may act as cell wall-binding domains, possibly targeting alginate (Michel et al. 2010).

One particularly interesting feature of the mannuronan C5-epimerase that was detected in ultrafiltrated SCM (encoded by the gene with LocusID Ec-20_004700) was the presence of three AGP backbone motifs. AGPs are large molecular weight glycoproteins (from 50 to more than 200 kDa) that were initially detected in land plant cell surfaces. Together with extensins and proline-rich proteins, AGPs belong to the group of hydroxyproline-rich-glycoproteins. AGPs are structurally characterized by a large carbohydrate moiety that accounts for 90-98% (w/w) of the molecule. Land plant AGPs consist predominantly of β -1,3-galactan main chains with β -1,6-galactan side chains, to which primarily arabinose and some other auxiliary sugar residues, such as rhamnose and glucuronic acid, are attached at their termini (Tan et al. 2012). This carbohydrate moiety is covalently linked via hydroxyproline to a relatively small protein backbone. AGPs have been shown to play crucial roles during development and reproduction in the land plant lineage (Hancock et al. 2005, Fu et al. 2007, Seifert and Roberts 2007). Developmental switches that appear to be influenced by AGPs in land plants include, for example, the acquisition of embryogenic capacity in *Daucus carota* (Kreuger and van Holst 1993, van Hengel et al. 2001), the role of the AGP-like protein xylogen in xylem formation in *Zinnia* and *Arabidopsis* (Motosue et al. 2004) and the role of AGP18 in the selection of viable megaspores in *Arabidopsis* (Demesa-Arévalo and Vielle-Calzada 2013). The phenotype associated with AGP18 is particularly interesting as this glycoprotein acts at the transition from the sporophyte to the gametophyte generation and loss of AGP18 leads to an arrest of gametogenesis (Acosta-García and Vielle-Calzada 2004). AGPs have also been shown to act as signal molecules during fertilisation. For example, tobacco TTS, which is produced by the stylar transmitting tissue, appears to act as a signalling factor to guide pollen tube growth (Cheung et al. 1995) and, in the land plant *Torenia fournieri*, ovular methyl-glucuronosyl arabinogalactan (AMOR) induces competency of the pollen tube to respond to ovular attractant peptides (Mizukami et al. 2016, Jiao et al. 2017).

In contrast to the situation for land plants, evidence for the presence of chimeric AGPs in brown algae has only been reported recently. Hervé *et al.* (2016) identified multiple genes in the *Ectocarpus* genome that were predicted to encode AGP-like core proteins. It is unknown if brown algal AGPs share conserved structural features with AGPs from land plants, however anti-AGP antibodies detect glycan epitopes in a variety of brown algae. The Hervé *et al.* (Hervé et al. 2016) study also presented evidence that AGP-like proteins are developmentally regulated during *Fucus* development and that developmental abnormalities occur if the action of the *Fucus* proteins is inhibited by the addition of AGP-reactive Yariv reagents.

In the current study, anti-AGP antibodies detected AGP glycan epitopes in SCM and addition of the Yariv reagent β -glucosyl to SCM led to an inhibition of its sporophyte-inducing activity. These observations suggest the interesting possibility that the sporophyte-inducing factor might be an AGP-like molecule. The lack of susceptibility of the sporophyte-inducing factor to proteinase K and heat treatment could be

interpreted as an argument against the factor corresponding to a protein but note that plant AGPs with similar heat and protease tolerances have been described. For example, xylogen has been shown to be heat resistant, although its activity is eliminated by proteases (Motose et al. 2001). AMOR was found to be both heat stable (100°C for 10 min) and maintained its activity following treatment with proteinase K. For the latter, a disaccharide, methyl-glucuronosyl galactose (4-Me-GlcA- β -(1/6)-Gal), was synthesised and shown to exhibit AMOR activity, suggesting that the protein moiety may not be important for the observed biological activity (Mizukami et al. 2016). By analogy, it is therefore possible that the active moiety of the *Ectocarpus* sporophyte-inducing factor is a saccharide molecule. Interestingly, SCM activity increased after storage for more than five weeks at 4°C (Fig. 1B) but no increase was observed when the medium was stored at -20°C for the same period of time (Fig. 1C). One possible interpretation of this observation might be that degradation of the diffusible factor led to the production of a more active degradation product such as a saccharide molecule.

In conclusion, this study has optimized conditions for the production of the sporophyte-inducing factor and has provided some initial information about its molecular nature that will be important for future attempts to purify the factor. Several different approaches provided evidence that the factor may be an AGP or an AGP-like molecule. Future work will be aimed at characterising the exact molecular nature of the factor and at investigating how the factor functions to induce the sporophyte program. On a broader scale, characterisation of the *Ectocarpus* factor may provide a starting point for the characterisation of analogous factors in other eukaryotic lineages. For example, moss sporophytes have been reported to produce a diffusible factor that induces apogamous sporophyte formation (Bauer, 1959) but the nature of this factor has not been investigated.

ACKNOWLEDGEMENTS

This work was supported by the Centre National de la Recherche Scientifique, the European Research Council (grant agreement 638240) and Sorbonne University. Support was also provided by the French National Research Agency via the investment expenditure programme Idealg (ANR-10-BTBR-04-02). HY was supported by a grant from the Chinese Scholarship Council (grant number 201608310119).

AUTHOR CONTRIBUTIONS

JMC and SMC conceived the study. HY carried out the experiments with input from DS for culture and bioassays and from MJ, CH and PP for biochemical and immunological analyses. LOC carried out the mass spectrometry analysis. HY analysed the data with input from DS, MJ, CH, PP, HY, SMC and JMC. HY, SMC and JMC wrote the first draft of the manuscript. All authors contributed to the final version. JMC and SMC contributed equally.

398

399 CONFLICT OF INTEREST

400 The authors have no conflict of interest.

401

402 References

- 403 Acosta-García, G. & Vielle-Calzada, J.-P. 2004. A classical arabinogalactan protein is essential for the
404 initiation of female gametogenesis in *Arabidopsis*. *Plant Cell*. 16:2614–28.
- 405 Arun, A., Coelho, S.M., Peters, A.F., Bourdareau, S., Pérès, L., Scornet, D., Strittmatter, M. et al. 2019.
406 Convergent recruitment of TALE homeodomain life cycle regulators to direct sporophyte
407 development in land plants and brown algae. *eLife*. 8:e43101.
- 408 Arun, A., Peters, N.T., Scornet, D., Peters, A.F., Cock, J.M. & Coelho, S.M. 2013. Non-cell autonomous
409 regulation of life cycle transitions in the model brown alga *Ectocarpus*. *New Phytol.* 197:503–10.
- 410 Bothwell, J.H., Marie, D., Peters, A.F., Cock, J.M. & Coelho, S.M. 2010. Role of endoreduplication and
411 apomeiosis during parthenogenetic reproduction in the model brown alga *Ectocarpus*. *New Phytol.*
412 188:111–21.
- 413 Brunelle, J. & Green, R. 2014. One-dimensional SDS-polyacrylamide gel electrophoresis (1D SDS-PAGE).
414 In *Methods in Enzymology*. Elsevier, pp. 151–9.
- 415 Cheung, A.Y., Wang, H. & Wu, H.M. 1995. A floral transmitting tissue-specific glycoprotein attracts pollen
416 tubes and stimulates their growth. *Cell*. 82:383–93.
- 417 Cock, J.M., Godfroy, O., Macaisne, N., Peters, A.F. & Coelho, S.M. 2014. Evolution and regulation of
418 complex life cycles: a brown algal perspective. *Curr Opin Plant Biol.* 17:1–6.
- 419 Coelho, S. & Cock, J. 2020. Brown algal model organisms. *Ann Rev Genet.* in press.
- 420 Coelho, S., Peters, A.F., Charrier, B., Destombe, C., Valero, M. & Cock, J. 2007. Complex life cycles of
421 multicellular eukaryotes: new approaches based on the use of model organisms. *Gene*. 406:152–70.
- 422 Coelho, S.M., Godfroy, O., Arun, A., Le Corguillé, G., Peters, A.F. & Cock, J.M. 2011. *OUROBOROS* is a
423 master regulator of the gametophyte to sporophyte life cycle transition in the brown alga *Ectocarpus*.
424 *Proc Natl Acad Sci USA*. 108:11518–23.
- 425 Coelho, S.M., Scornet, D., Rousvoal, S., Peters, N.T., Darteville, L., Peters, A.F. & Cock, J.M. 2012. How to
426 cultivate *Ectocarpus*. *Cold Spring Harb Protoc.* 2012:258–61.
- 427 Demesa-Arévalo, E. & Vielle-Calzada, J.-P. 2013. The classical arabinogalactan protein AGP18 mediates
428 megaspore selection in *Arabidopsis*. *Plant Cell*. 25:1274–87.
- 429 Dittami, S.M., Corre, E., Brillet-Guéguen, L., Lipinska, A.P., Pontoizeau, N., Aite, M., Avia, K. et al. 2020.
430 The genome of *Ectocarpus subulatus* - A highly stress-tolerant brown alga. *Mar. Genomics*. 100740.
- 431 Dupres, V., Alsteens, D., Wilk, S., Hansen, B., Heinisch, J.J. & Dufrêne, Y.F. 2009. The yeast Wsc1 cell
432 surface sensor behaves like a nanospring *in vivo*. *Nat. Chem. Biol.* 5:857–62.
- 433 Fischl, R., Bertelsen, K., Gaillard, F., Coelho, S., Michel, G., Klinger, M., Boyen, C. et al. 2016. The cell-wall
434 active mannuronan C5-epimerases in the model brown alga *Ectocarpus*: From gene context to
435 recombinant protein. *Glycobiology*. 26:973–83.
- 436 Fu, H., Yadav, M.P. & Nothnagel, E.A. 2007. *Physcomitrella patens* arabinogalactan proteins contain
437 abundant terminal 3-O-methyl-L: -rhamnosyl residues not found in angiosperms. *Planta*. 226:1511–
438 24.

439 Gane, A.M., Craik, D., Munro, S.L., Howlett, G.J., Clarke, A.E. & Bacic, A. 1995. Structural analysis of the
 440 carbohydrate moiety of arabinogalactan-proteins from stigmas and styles of *Nicotiana glauca*.
 441 *Carbohydr. Res.* 277:67–85.

442 Hancock, C.N., Kent, L. & McClure, B.A. 2005. The stylar 120 kDa glycoprotein is required for S-specific
 443 pollen rejection in *Nicotiana*. *Plant J. Cell Mol. Biol.* 43:716–23.

444 Hervé, C., Siméon, A., Jam, M., Cassin, A., Johnson, K.L., Salmeán, A.A., Willats, W.G.T. et al. 2016.
 445 Arabinogalactan proteins have deep roots in eukaryotes: identification of genes and epitopes in brown
 446 algae and their role in *Fucus serratus* embryo development. *New Phytol.* 209:1428–41.

447 Jiao, J., Mizukami, A.G., Sankaranarayanan, S., Yamguchi, J., Itami, K. & Higashiyama, T. 2017. Structure-
 448 Activity Relation of AMOR Sugar Molecule That Activates Pollen-Tubes for Ovular Guidance. *Plant*
 449 *Physiol.* 173:354–63.

450 Johnson, K.L., Jones, B.J., Bacic, A. & Schultz, C.J. 2003. The fasciclin-like arabinogalactan proteins of
 451 *Arabidopsis*. A multigene family of putative cell adhesion molecules. *Plant Physiol.* 133:1911–25.

452 Kreuger, M. & van Holst, G.-J. 1993. Arabinogalactan proteins are essential in somatic embryogenesis of
 453 *Daucus carota* L. *Planta*. 189:243–8.

454 Lodder, A.L., Lee, T.K. & Ballester, R. 1999. Characterization of the Wsc1 protein, a putative receptor in the
 455 stress response of *Saccharomyces cerevisiae*. *Genetics*. 152:1487–99.

456 Lommel, M., Bagnat, M. & Strahl, S. 2004. Aberrant processing of the WSC family and Mid2p cell surface
 457 sensors results in cell death of *Saccharomyces cerevisiae* O-mannosylation mutants. *Mol. Cell. Biol.*
 458 24:46–57.

459 Michel, G., Tonon, T., Scornet, D., Cock, J.M. & Kloareg, B. 2010. The cell wall polysaccharide metabolism
 460 of the brown alga *Ectocarpus siliculosus*. Insights into the evolution of extracellular matrix
 461 polysaccharides in Eukaryotes. *New Phytol.* 188:82–97.

462 Mizukami, A.G., Inatsugi, R., Jiao, J., Kotake, T., Kuwata, K., Ootani, K., Okuda, S. et al. 2016. The AMOR
 463 Arabinogalactan Sugar Chain Induces Pollen-Tube Competency to Respond to Ovular Guidance.
 464 *Curr. Biol. CB*. 26:1091–7.

465 Motose, H., Sugiyama, M. & Fukuda, H. 2001. An arabinogalactan protein(s) is a key component of a fraction
 466 that mediates local intercellular communication involved in tracheary element differentiation of zinnia
 467 mesophyll cells. *Plant Cell Physiol.* 42:129–37.

468 Motose, H., Sugiyama, M. & Fukuda, H. 2004. A proteoglycan mediates inductive interaction during plant
 469 vascular development. *Nature*. 429:873–8.

470 Müller, D.G. 1967. Generationswechsel, Kernphasenwechsel und Sexualität der Braunalge *Ectocarpus*
 471 *siliculosus* im Kulturversuch. *Planta*. 75:39–54.

472 Nyvall, P., Corre, E., Boisset, C., Barbeyron, T., Rousvoal, S., Scornet, D., Kloareg, B. et al. 2003.
 473 Characterization of mannuronan C-5-epimerase genes from the brown alga *Laminaria digitata*. *Plant*
 474 *Physiol.* 133:726–35.

475 Ohsawa, S., Yurimoto, H. & Sakai, Y. 2017. Novel function of Wsc proteins as a methanol-sensing machinery
 476 in the yeast *Pichia pastoris*. *Mol. Microbiol.* 104:349–63.

477 Oide, S., Tanaka, Y., Watanabe, A. & Inui, M. 2019. Carbohydrate-binding property of a cell wall integrity
 478 and stress response component (WSC) domain of an alcohol oxidase from the rice blast pathogen
 479 *Piricularia oryzae*. *Enzyme Microb. Technol.* 125:13–20.

480 Paulsen, B.S., Craik, D.J., Dunstan, D.E., Stone, B.A. & Bacic, A. 2014. The Yariv reagent: behaviour in
 481 different solvents and interaction with a gum arabic arabinogalactan-protein. *Carbohydr. Polym.*
 482 106:460–8.

483 Peters, A.F., Marie, D., Scornet, D., Kloareg, B. & Cock, J.M. 2004. Proposal of *Ectocarpus siliculosus*
484 (Ectocarpales, Phaeophyceae) as a model organism for brown algal genetics and genomics. *J Phycol.*
485 40:1079–88.

486 Peters, A.F., Scornet, D., Ratin, M., Charrier, B., Monnier, A., Merrien, Y., Corre, E. et al. 2008. Life-cycle-
487 generation-specific developmental processes are modified in the *immediate upright* mutant of the
488 brown alga *Ectocarpus siliculosus*. *Development*. 135:1503–12.

489 Popper, Z.A., Michel, G., Hervé, C., Domozych, D.S., Willats, W.G.T., Tuohy, M.G., Kloareg, B. et al. 2011.
490 Evolution and diversity of plant cell walls: from algae to flowering plants. *Ann Rev Pl Biol*. 62:567–
491 90.

492 Seifert, G.J. & Roberts, K. 2007. The biology of arabinogalactan proteins. *Annu. Rev. Plant Biol.* 58:137–61.

493 Tan, L., Showalter, A.M., Egelund, J., Hernandez-Sanchez, A., Doblin, M.S. & Bacic, A. 2012.
494 Arabinogalactan-proteins and the research challenges for these enigmatic plant cell surface
495 proteoglycans. *Front. Plant Sci.* 3:140.

496 van Hengel, A.J., Tadesse, Z., Immerzeel, P., Schols, H., van Kammen, A. & de Vries, S.C. 2001. N-
497 acetylglucosamine and glucosamine-containing arabinogalactan proteins control somatic
498 embryogenesis. *Plant Physiol.* 125:1880–90.

499 Wawra, S., Fesel, P., Widmer, H., Neumann, U., Lahrmann, U., Becker, S., Hehemann, J.-H. et al. 2019. FGB1
500 and WSC3 are in planta-induced β -glucan-binding fungal lectins with different functions. *New Phytol.*
501 222:1493–506.

502 Yao, H., Scornet, D., Badis, Y., Peters, A.F., Jam, M., Hervé, C., Potin, P. et al. 2020. Production and Bioassay
503 of a Diffusible Factor that Induces Gametophyte-to-Sporophyte Developmental Reprogramming in
504 the Brown Alga *Ectocarpus*. *Bio-Protoc.* 10:e3753–e3753.

507 **TABLE**

508 Table 1. Proteins detected in concentrated sporophyte-conditioned medium following SDS-PAGE, excision
 509 of bands >50 kDa and mass spectrometry. SP, signal peptide; TM, transmembrane domain; WSC, cell-wall
 510 integrity and stress-response component 1 domain; MC5E, mannuronan C-5-epimerase; AGP,
 511 arabinogalactan protein; G/SdH, glucose/sorbose dehydrogenase; PLF/V, pectin lyase fold/virulence
 512 factor; PS8/S53, petidase S8/S53; HIP, high-CO₂ inducible periplasmic; PKinase, protein kinase; RCC1,
 513 regulator of chromosome condensation; SBBP, six-bladed beta-propeller, TolB-like; Gal, galactose-binding;
 514 AP, alkaline-phosphatase-like; SGNH, SGNH hydrolase-type esterase; HLRR, hypothetical leucine-rich
 515 repeat; vWA, von Willebrand factor type A; FA58C, Coagulation factor 5/8 C-terminal type domain; HP,
 516 Haem peroxidase; CA, carbonic anhydrase; GH114, N-terminal glycosyl-hydrolase-114-associated domain.

Description	Locus ID	Domains	Major Component
Mannuronan C-5-epimerases (short name indicated)	Ec-20_004700 (MEP4)	SP, 2xWSC, MC5E, AGP	No
	Ec-00_006370 (MEP5)	SP, MC5E	Yes
	Ec-00_006380 (MEP6)	SP, MC5E	Yes
	Ec-11_000400 (MEP2)	SP, MC5E	Yes
	Ec-27_006700 (MEP28)	SP, MC5E	Yes
	Ec-18_000130 (MEP7)	MC5E	Yes
	Ec-00_010800 (MEP13)	TM, 2xWSC, MC5E	Yes
	Ec-18_000150 (MEP8)	SP, MC5E	No
WSC domain protein	Ec-17_000970	SP, WSC	Yes
Glucose/Sorbose dehydrogenase and WSC domain protein	Ec-00_007780	SP, G/SdH, 5xWSC	Yes
	Ec-09_000520	SP, G/SdH, WSC	Yes
	Ec-13_003060	SP, G/SdH, 3xWSC	No
High-CO ₂ inducible periplasmic domain protein	Ec-26_006220	SP, HIP	Yes
Pectin lyase fold proteins	Ec-03_002910	SP, PLF/V	No
	Ec-22_003520	SP, PLF/V	Yes
Seven-bladed beta-propeller domain receptor kinase	Ec-22_003740	SP, RCC1, TM, PKinase	Yes
TolB-like six-bladed beta-propeller proteins	Ec-15_001700	SP, SBBP	Yes
	Ec-02_001340	SP, SBBP	No

Galactose-binding domain-like proteins	Ec-21_001340	SP, 3xGal, PLF/V	No
	Ec-21_001320	SP, Gal, PLF/V	No
	Ec-21_001360	SP, 3xGal, PLF/V	Yes
	Ec-21_001350	SP, 2xGal, PLF/V	Yes
Seven-bladed beta-propeller domain protein	Ec-16_004480	SP, RCC1, TM	Yes
Secreted protein similar to EsV-1-163	Ec-26_002660	SP	No
	Ec-26_005190	SP	No
asn/thr-rich large protein family protein	Ec-02_004700	SP	No
Secreted alkaline phosphatase	Ec-01_007880	SP, AP	Yes
KP-43 peptidase	Ec-06_007900	PS8/S53, Gal	No
SGNH hydrolase-type esterase domain	Ec-09_000420	SP, SGNH	No
Hypothetical leucine rich repeat protein	Ec-09_004400	SP, Gal, FA58C, LRR	No
Von Willebrand factor type A domain protein	Ec-21_003770	vWA	No
Catalase (imm downregulated 7)	Ec-26_000310	SP, HP	Yes
FAS1 domain protein	Ec-27_005070	FAS1, TM	Yes
carbonic anhydrase alpha type	Ec-27_005680	SP, CA	No
conserved unknown protein	Ec-05_001620	SP	Yes
GH114 glycosyl-hydrolase	Ec-28_003870	GH114	Yes

517

518

521
522

523 Fig. 1. Effect of different parameters on the activity of the diffusible factor. (A) Effect of unilocular
524 sporangium release time on uni-spore susceptibility to the diffusible sporophyte-inducing factor. Percent of
525 germlings that exhibited sporophyte morphology following the release of uni-spores from unilocular
526 sporangia 12-24 h, 24-48 h or >48 h after dissection and transfer to either SCM or to PES as a negative
527 control. (B,C) The sporophyte-inducing factor is stable when SCM is stored at 4°C or -20°C. (B) Bioassay

528 activity after storage at 4°C. (C) Bioassay activity after storage at -20°C. (D,E) Effect of sporophyte culture
529 conditions on production of the sporophyte-inducing diffusible factor. (D) Effect of time in culture, (E)
530 Effect of light intensity. Error bars indicate standard error of the mean, letters above bars indicate significant
531 differences (Wilcoxon-Mann-Whitney test, p -value <0.05; see table S1 for details of statistical tests). PES,
532 Provasoli-enriched natural seawater; LL, low light; NL, normal light; HL, high light; n, number of replicates;
533 i, number of individual germlings counted.

534

535
536

537
538

539 Fig. 2. Sporophyte-inducing activity of different SCM size fractions. (A) Size fractionation of SCM using
540 ultrafiltration. The diagrams indicate input, flow through and retentate volumes. Washes with deionised
541 water were carried out after completion of SCM filtration and the retentate volumes correspond to the final

542 volumes after washing where relevant. The degree to which a sample was concentrated following
543 ultrafiltration is indicated by the number in brackets under the text indicating the molecular sizes of
544 molecules in the recovered samples, e.g. (5x), 10-50 kDa molecules were concentrated 5-fold compared to
545 the initial SCM preparation before filtration. (B) SCM size fractions, separated by ultrafiltration, were tested
546 on uni-spores for sporophyte-inducing activity. All samples were diluted to a concentration equivalent to 1X
547 SCM before carrying out the bioassays. Error bars indicate standard error of the mean, letters above bars
548 indicate significant differences (Wilcoxon-Mann-Whitney test, p -value <0.05 ; see table S2 for details of
549 statistical tests). PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; n,
550 number of replicates; i, number of individual germlings counted.

551

Fig. 3. Identification of candidate proteins for the diffusible sporophyte-inducing factor by mass spectroscopy analysis of protein bands detected in the >30 kDa fraction of concentrated SCM. (A) Coomassie stained SDS-PAGE gel showing the protein bands detected in concentrated SCM. Numbers and bars on the right indicate the fragments that were excised for mass spectrometry analysis. (B) Domain structure of the Ec-20_004700 protein. Green boxes indicate AGP protein cores with putative attached glycan groups. The Ec-20_004700 protein was detected in band 4.

563
564

565
566

567 Fig. 4. Effect of heat and proteinase K treatment on the sporophyte-inducing factor. SCM was autoclaved at
568 121°C for 20 min (right panel) or incubated with proteinase K (100 µg/ml) at 37°C for 1h followed by
569 autoclaving at 121°C for 20 min (left panel). Error bars indicate standard error of the mean, letters above
570 bars indicate significant differences (Wilcoxon-Mann-Whitney test, *p*-value <0.05; see table S3 for details of
571 the statistical tests). PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; pK-
572 SCM, proteinase K and heat treated SCM; heat-SCM, heat treated SCM; n, number of replicates; i, number
573 of individual germlings counted.

574

575
576

577
578

579 Fig. 5. Detection of AGP epitopes in sporophyte-conditioned medium. Samples were serially diluted in five
580 steps of five-fold dilution and 1 μl of each dilution spotted for testing with the anti-AGP monoclonal
581 antibodies JIM4, JIM8, JIM13, JIM16, LM2, LM14 and MAC207. Gum arabic (starting concentration 1
582 mg/ml) was used as a positive control. Molecules >50 kDa were 400-fold more concentrated in the neat uf-
583 SCM sample compared to the neat SCM sample. GA, gum arabic; SCM, sporophyte-conditioned medium;
584 uf-SCM, ultrafiltrated SCM; control, no primary antibody.

585

586
587

588
589
590
591
592
593
594
595
596
597

Fig. 6. Inhibition of the sporophyte-inducing activity of SCM following addition of a Yariv reagent. SCM was incubated with either β -D-glucosyl Yariv reagent or with β -D-mannosyl Yariv reagent as a control. Error bars indicate standard error of the mean, letters above bars indicate significant differences (Wilcoxon-Mann-Whitney test, p -value <0.05 ; see table S4 for details of the statistical tests). PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; β Glc-Y, β -D-glucosyl Yariv reagent; β Man-Y, β -D-mannosyl Yariv reagent; n, number of replicates; i, number of individual germlings counted.

SUPPLEMENTARY TABLES

Table S1. Pairwise Wilcoxon-Mann-Whitney tests comparing the levels of conversion to the sporophyte generation by the diffusible factor under different experimental conditions. Refers to Fig. 1. PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; LL, low light; NL, normal light; HL, high light.

comparison	<i>W</i>	adjusted <i>p</i> -value
Fig. 1A		
PES (<24h) / SCM (<24h)	399.5	0.0154
PES (<24h) / PES (24-48h)	679	0.0170
PES (<24h) / SCM (24-48hs)	816	1.0000
PES (<24h) / PES (>48hc)	707.5	0.0273
PES (<24h) / SCM (>48hs)	568.5	1.0000
SCM (<24h) / PES (24-48hc)	1180.5	2.0e ⁻⁰⁶
SCM (<24h) / SCM (24-48hs)	1614.5	0.1277
SCM (<24h) / PES (>48hc)	1241.5	2.3e ⁻⁰⁵
SCM (<24h) / SCM (>48hs)	1120	0.0313
PES (24-48h) / SCM (24-48hs)	607	0.0027
PES (24-48h) / PES (>48hc)	563	1.0000
PES (24-48h) / SCM >48hs	427	0.0166
SCM (24-48hs) / PES (>48hc)	1402	0.0042
SCM (24-48hs) / SCM >48hs	1206.5	1.0000
PES (>48hc) / SCM (>48hs)	466.5	0.0
Fig. 1B		
PES / 0 weeks	282	5.1 e ⁻¹¹
PES / 1 weeks	41	6.1e ⁻⁰⁷
PES / 3 weeks	437	1.7e ⁻¹¹
PES / 5 weeks	63.5	2.6e ⁻¹³
PES / 8 weeks	2	3.5 e ⁻⁰⁶
0 weeks / 1 weeks	138	1.0000
0 weeks / 3 weeks	482	1.0000
0 weeks / 5 weeks	145	0.0326
0 weeks / 8 weeks	32.5	0.1234
1 weeks / 3 weeks	133.5	1.0000
1 weeks / 5 weeks	29	0.0289
1 weeks / 8 weeks	4	0.0541
3 weeks / 5 weeks	162.5	0.0052
3 weeks / 8 weeks	40.5	0.1068
5 weeks / 8 weeks	64.5	1.0000
Fig. 1C		
PES / 0 weeks	954.5	1.3e ⁻¹¹

PES / 1 weeks	316	4.0e ⁻⁰⁷
PES / 3 weeks	326	1.2e ⁻⁰⁷
PES / 5 weeks	241	8.8e ⁻¹²
PES / 8 weeks	573.5	0.0002
0 weeks / 1 weeks	1086	1.0000
0 weeks / 3 weeks	1196	1.0000
0 weeks / 5 weeks	1394	1.0000
0 weeks / 8 weeks	1577.5	0.0346
1 weeks / 3 weeks	409.5	1.0000
1 weeks / 5 weeks	431	1.0000
1 weeks / 8 weeks	586	0.0705
3 weeks / 5 weeks	457.5	1.0000
3 weeks / 8 weeks	605	0.1484
5 weeks / 8 weeks	830	0.0061
Fig. 1D		
PES / 4 weeks	1116.5	0.00221
PES / 6 weeks	195.5	6.6e ⁻¹¹
PES / 8 weeks	131.5	2.6e ⁻¹²
PES / 10 weeks	1058.5	0.42732
4 weeks / 6 weeks	190	0.00473
4 weeks / 8 weeks	142.5	0.00040
4 weeks / 10 weeks	511.5	0.32191
6 weeks / 8 weeks	173	0.32191
6 weeks / 10 weeks	434	0.00066
8 weeks / 10 weeks	456.5	0.00011
Fig. 1E		
PES / LL	938	1.501e ⁻⁰⁵
PES / NL	610.5	0.000
PES / HL	233.5	1.134e ⁻⁰⁵
LL / NL	1461	0.429
LL / HL	785.5	0.166
NL / HL	571.5	0.662

605

606

607

608

609

Table S2. Pairwise Wilcoxon-Mann-Whitney tests comparing the levels of conversion to the sporophyte generation induced by treatment with different SCM size fractions. Refers to Fig. 2. PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; kDa, kiloDaltons.

comparison	<i>W</i>	adjusted <i>p</i> -value
Fig. 2		
PES / SCM	14	1.5 e ⁻⁰⁵
PES / <100 kDa	45	0.00032
PES / >50 kDa	0	3.5e ⁻⁰⁷
PES / 10-50 kDa	202	1.0000

SCM / <100 kDa	208	1.0000
SCM / >50 kDa	209	1.0000
SCM / 10-50 kDa	354	1.0 e ⁻⁰⁵
<100 kDa / >50 kDa	197.5	1.0000
<100 kDa / 10-50 kDa	362.5	0.00018
>50 kDa / 10-50 kDa	614	4.3e ⁻⁰⁷

Table S3. Pairwise Wilcoxon-Mann-Whitney tests comparing the effects of heat and proteinase K treatments on levels of conversion to the sporophyte generation by the diffusible factor. Refers to Fig. 4. PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; pk, proteinase K.

comparison	<i>W</i>	adjusted <i>p</i> -value
Fig. 4		
PES / SCM	165.5	5.1e ⁻¹¹
PES / heat-SCM	86.5	3.7e ⁻¹⁰
SCM / heat-SCM	3385.5	0.1
PES / SCM	0	0.00059
PES / pk-SCM	0	0.00013
SCM / pk-SCM	45.5	0.03785

Table S4. Pairwise Wilcoxon-Mann-Whitney tests comparing the effects of treatments with different Yariv reagents on levels of conversion to the sporophyte generation by the diffusible factor. Refers to Fig. 6. PES, Provasoli-enriched natural seawater; SCM, sporophyte-conditioned medium; βGlc-Y, β-D-glucosyl Yariv reagent; βMan-Y, β-D-mannosyl Yariv reagent.

comparison	<i>W</i>	adjusted <i>p</i> -value
Fig. 6		
PES / SCM	1.5	2.4 e ⁻⁰⁵
PES / SCM+βMan-Y	44.5	0.0016
PES / SCM+βGlc-Y	44.5	0.0075
SCM / SCM+βMan-Y	111	0.5305
SCM / SCM+βGlc-Y	115	0.0369
SCM+βMan-Y / SCM+βGlc-Y	127.5	0.5305