



# Ultrastructural research of spermiogenesis in two sponges, *Crellomima imparidens* and *Hymedesmia irregularis* (Demospongiae): new evidence of sperms with acrosome in sponges

Alexander Ereskovsky, Daria Tokina

## ► To cite this version:

Alexander Ereskovsky, Daria Tokina. Ultrastructural research of spermiogenesis in two sponges, *Crellomima imparidens* and *Hymedesmia irregularis* (Demospongiae): new evidence of sperms with acrosome in sponges. *Journal of Morphology*, 2022, 283 (3), pp.333-345. 10.1002/jmor.21446 . hal-03527256

**HAL Id: hal-03527256**

**<https://cnrs.hal.science/hal-03527256>**

Submitted on 26 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1                    **Ultrastructural research of spermiogenesis in two sponges, *Crellomima***  
2 ***imparidens* and *Hymedesmia irregularis* (Demospongiae): new evidence of sperms with**  
3 **acrosome in sponges**

4  
5                    **Research Highlights:**

- 6    1. The ultrastructure of sperm were studied in two marine demosponges, *Crellomima*  
7 *imparidens* and *Hymedesmia irregularis* (order Poecilosclerida)  
8    2. The bundles of microtubules arranged along the nucleus during spermiogenesis found in  
9 both species.  
10    3. Sperm nucleus have elongated and helical shape and twisted chromatin.  
11    4. The spermatozoon of both species has an acrosome at apical part.  
12  
13

14                    Alexander Ereskovsky <sup>1,2,3\*</sup>, Daria Tokina<sup>2</sup>

15    <sup>1</sup>13007, France, Marseille, Institut Méditerranéen de Biodiversité et d'Ecologie Marine et  
16 Continentale (IMBE), Aix Marseille University, CNRS, IRD, Avignon University, Station  
17 Marine d'Endoume, Rue de la Batterie des Lions.

18    <sup>2</sup>199034, Russia, Saint-Petersburg Department of Embryology, Faculty of Biology, Saint-  
19 Petersburg State University, Universitetskaya nab. 7/9.

20    <sup>3</sup>119334, Russia, Moscow, Koltzov Institute of Developmental Biology of Russian  
21 Academy of Sciences, Vavilova street 26.  
22

23                    **Abstract**

24                    Details of spermatogenesis and sperm organization are often useful for reconstructing  
25 the phylogeny of closely related taxa of invertebrates. Here, the spermiogenesis and the  
26 ultrastructure of sperm were studied in two marine demosponges, *Crellomima imparidens*  
27 and *Hymedesmia irregularis* (order Poecilosclerida). In *C. imparidens* and *H. irregularis*  
28 we found bundles of microtubules arranged along the nucleus during spermiogenesis.  
29 These bundles derived from the basal body of axoneme, reaching the apical pole of the  
30 cell. In *C. imparidens* the microtubules surround the nucleus, forming the manchette. In *H.*  
31 *irregularis* the microtubules pass along only one side of the cell periphery. During  
32 spermiogenesis, the nucleus stretches and elongates. In both species the nucleus is twisted  
33 into a spiral structure. We suppose that the manchette of microtubules could be responsible  
34 for controlling the elongation and shaping of the sperm nucleus to a helical form and for  
35 the twisting and/or condensation of chromatin in these sponges. The spermatozoon of both  
36 species has an elongated shape. Its apical part has an acrosome, which is dome-shaped in  
37 *C. imparidens* and flattened and lenticular in *H. irregularis*. The cytoplasm of the  
38 spermatozoa contains some small mitochondria, and proximal and distal centrioles

1 arranged at an angle to each other. There is a small volume of residual cytoplasm with dark  
2 glycogen-like granules. The axoneme of the spermatid and the flagellum of the sperm of  
3 both sponges is located in the deep tunnel-like cytoplasmic depression. The comparison of  
4 spermatozoa morphology of different species of the order Poecilosclerida demonstrates  
5 that the knowledge of variation within genera and families can give valuable insights into  
6 the significance of many characters proposed for phylogenetic studies of this order.

7  
8 **Keywords:** spermiogenesis, sponges, acrosome, ultrastructure

## 9 10 **1. Introduction**

11  
12 Electron microscopic observations of gametogenesis in sponges (Porifera) have  
13 demonstrated a high morphological diversity of spermatozoa in these animals. Moreover,  
14 all three morphological groups of metazoan sperm cells can be found in sponges from  
15 different taxa: “primitive spermatozoa,” “derived or modified spermatozoa,” and “aberrant  
16 spermatozoa” according to Reunov (2005). Accordingly, there are sponge species  
17 possessing sperm with an acrosome, which was earlier considered as a trait of more highly  
18 organized groups of animals (Afzelius, 1978; Berruti et al., 1978; Baccetti, 1984). The  
19 presence of the acrosome in spermatozoa has been shown in the class Homoscleromorpha  
20 (Gaino et al., 1986; Baccetti et al., 1986; Boury-Esnaul & Jamieson, 1999; Ereskovsky,  
21 2010; Riesgo et al., 2007a), in one species of the class Calcarea (Nakamura et al., 1998)  
22 and, finally, in some Demospongiae from the order Poecilosclerida: *Crambe crambe*  
23 (Tripepi et al., 1984; Riesgo & Maldonado, 2009), *Lycopodina occidentalis* and *L.*  
24 *hypogea* (Riesgo et al., 2007b; Ereskovsky et al. In prep.).

25 Ultrastructural studies of spermiogenesis and sperm morphology in different  
26 invertebrates have provided many characters used as apomorphies as well as  
27 morphological evidence used in taxonomy and phylogeny (for review see Jamieson et al.,  
28 1995). Differences in sperm morphology and spermiogenesis have been found at high  
29 taxonomic levels within various animal groups such as Turbellaria (Watson & Rhode,  
30 1995), Cestoda (Ba & Marchand, 1995), Gastrotricha (Ferraguti & Balsamo, 1995),  
31 Polychaeta (Jamieson & Rouse, 1989), Cnidaris (Hinsch, 1974), and Polyplacophora  
32 (Buckland-Nicks, 2006).

33 However, sperm morphology has never been used in taxonomy and phylogeny of  
34 sponges, mostly because spermatogenesis has been studied at the ultrastructural level only

in a few species from different sponge taxa. Most studies of spermatogenesis in sponges have been made on representatives of their largest class, Demospongiae. Spermatogenesis has been examined using TEM in 27 species from 10 orders of this class (Table 1). The order Poecilosclerida has been studied best, with information on the ultrastructure of spermatids or spermatozoa being available for 9 species (Table 1) (Efremova et al., 1987; De Vos et al., 1990; Riesgo et al., 2007b; Riesgo & Maldonado, 2009; Pérez-Porro et al., 2012; Vasconcellos et al., 2019; Ereskovsky et al., In prep).

The aim of this work was to perform an electron microscopic study of the late stages of spermatogenesis—spermiogenesis and sperm formation—in two sponge species from different families of the order Poecilosclerida: *Crellomima imparidens* and *Hymedesmia irregularis*.

## 2. Materials and methods

Sponges *Crellomima imparidens* Rezvoj, 1925 (family Crellidae) and *Hymedesmia irregularis* Lundbeck, 1910 (family Hymedesmiidae) (Fig. 1 a,b) were collected by SCUBA diving from June to August 2006 and from August to September 2009 in the Chupa Inlet located in the innermost part of the Kandalaksha Bay, the White Sea, at depths of 7-12 m. Pieces of each specimen were fixed immediately after collection. For electron microscopy the sponges were cut into cubes of about 1mm. These cubes were prefixed in 1% OsO<sub>4</sub> (Spi Supplies, West Chester) for 10 min and fixed in 2.5 % glutaraldehyde (Ted Pella, Redding) in phosphate buffer at room temperature for 1 h. After fixation, sponge samples were washed in the phosphate buffer and postfixated in 1% OsO<sub>4</sub> in phosphate buffer for 1 h. After postfixation, the samples were kept in 5% HF solution for 3 hr at 4°C in order to dissolve the spicules. Samples were dehydrated through a graded ethanol series at RT and stored in 70% ethanol at 4°C.

For semithin sections and transmission electron microscopy, the specimens were embedded in Araldite (Sigma-Aldrich) according to the manufacturer's instructions. Semithin sections (1 µm) were cut on an Ultramicrotome PowerTome XL (RMC Boeckeler, Tucson), and then stained with 1% toluidine blue–0.2% methylene blue mixture. They were studied under a Leica DM5000B (Leica) and a Wild M20 microscope (Wild). Digital photos were taken with a Leica DMLB microscope (Leica) using the Evolution LC color photo capture system (Media Cybernetics, Rockville).

1 Ultrathin sections (60–80 nm) were cut with an Ultramicrotome PowerTome XL,  
2 equipped with a Drukkert 45° diamond knife and contrasted with 4% aqueous uranyl  
3 acetate and 0.4% lead citrate according to Reynolds (1963). Ultrathin sections were studied  
4 under Jeol JEM-100B (Jeol, Akishima, Japan) transmission electron microscope.

### 5 6 **3. Results**

#### 7 8 **3.1. Sperm cysts**

9 Spermatogenesis in both *Crellomima imparidens* and *Hymedesmia irregularis* occurs  
10 in sperm cysts, temporary oval structures with a diameter of about 38–42  $\mu\text{m}$  in *C.*  
11 *imparidens* and 26–40 x 17–22  $\mu\text{m}$  in *H. irregularis* (Fig. 2 a–d). Spermatogenesis within a  
12 cyst is relatively synchronous. However, in different cysts of the same sponge  
13 spermatocytes can be at different stages of spermatogenesis. The cyst wall is single-  
14 layered, formed with flat follicular cells with long inwardly directed pseudopodia (Fig. 2  
15 c). The cyst cells have a large spherical anucleolate nucleus (about 3.8  $\mu\text{m}$  in diameter in  
16 *C. imparidens* and 4  $\mu\text{m}$  in *H. irregularis*) with islands of heterochromatin. The cytoplasm  
17 contains numerous spherical heterophagosomes (from 0.5 to 2.5  $\mu\text{m}$ ) and electron  
18 transparent vacuoles of different diameter. The cyst cells contact by simply overlying each  
19 other.

#### 20 21 **3.2. Spermiogenesis in *Crellomima imparidens***

22 Early spermatids are oval cells with a diameter about 2.9  $\mu\text{m}$  in cross section (Figs. 2a,  
23 3a). Cells in groups of four are often interconnected by cytoplasmic bridges (Fig. 3b).  
24 Spermatids have a flagellum. The nuclei of early spermatids are oval with a diameter of  
25 about 1.8  $\mu\text{m}$ . In the cytoplasm there are rounded mitochondria (diameter about 1.8  $\mu\text{m}$ )  
26 and the Golgi apparatus, represented by flat cisterns located near the nucleus. These  
27 cisterns remain in the residual cytoplasm in the course of sperm maturation. During  
28 maturation chromatin in the nuclei of spermatids condenses. This process begins at the  
29 periphery, while the central part of the nucleus remains lighter. The nuclei gradually  
30 acquire an elongated cylindrical shape. The spermatids themselves also elongate, reaching  
31 up to 6.3  $\mu\text{m}$  in length. The residual cytoplasm accumulates in the basal part of spermatids,  
32 which is discharged at the end of spermiogenesis.

1 As the sperm stretches, the nucleus is twisted into a spiral structure (Figs. 3 c,d; Fig. 4  
2 a ; 5– scheme long.). The sperm nucleus is electron-dense, homogeneous, 4.5  $\mu\text{m}$  long. It is  
3 narrower in the proximal part than in the distal part (diameter, respectively, 0.5 and 0.23  
4  $\mu\text{m}$ ). In the central part of the distal pole, the nucleus has a small funnel-shaped depression  
5 up to 0.06 - 0.11  $\mu\text{m}$  deep (Figs. 3 e, m-o; Fig. 4 a). Microtubules are uniformly arranged  
6 around the nucleus, in parallel with it (Figs. 3e, f, g; 4 a, c-e; 5-scheme section). They  
7 begin from the basal body of the axoneme (Fig. 3 e) and reach the apical part of the  
8 nucleus.

9 There is one flagellum in the basal part of the sperm. It has a typical eukaryotic  
10 axoneme (9 +2), which is located in the cytoplasmic depression with a depth of 1.1  $\mu\text{m}$   
11 (Fig. 3 m). The basal body lies at the base of the axoneme (Figs. 3 n, o; 4a). An accessory  
12 centriole is located at an angle to the basal body (Fig. 3 n). A short rootlet arises from the  
13 terminal part of the basal body (Fig. 3 n) and extends to contact the basal part of the  
14 nucleus.

15 In general, the sperm cytoplasm is fairly homogeneous. There are small mitochondria  
16 and lipid droplets in its basal part and several cisterns derived from the Golgi apparatus  
17 around the axoneme.

18 The acrosome forms in early spermatids. It is located in the immediate vicinity of the  
19 Golgi apparatus. In the early stages, it is rounded, with a diameter of about 1.2  $\mu\text{m}$ . In  
20 more mature spermatids, the acrosome occupies an apical position, adjacent to the distal  
21 part of the nucleus. It is dome-shaped, homogenous, with a diameter of 0.36  $\mu\text{m}$  and a  
22 height of 0.31  $\mu\text{m}$  (Figs. 3 h-l, 4 a, b). However, it has two electron-transparent regions: a  
23 lenticular region in the apical part and an annular groove in the middle part (Fig. 3 h-j).  
24 Moreover, its basal part, which is adjacent to the nucleus, is filled with a darker electron-  
25 dense material, while the opposite part, which faces the cytoplasmic membrane, is more  
26 electron-transparent (Fig. 3 h-j). The acrosome has its own membrane.

27 Sperm are released from the cyst due to the disruption of cellular contacts of the sperm  
28 cyst.

### 30 **3.3. Spermiogenesis in *Hymedesmia irregularis***

31 Early spermatids of *Hymedesmia irregularis* are oval monoflagellated cells, with an  
32 elongated nucleus, the chromatin of which begins to condense (Figs. 2b; 6 a). Their  
33 cytoplasm contains numerous small mitochondria, small electron-transparent vacuoles,  
34 lipid droplets and several elongated, sinuous cisterns derived from the Golgi apparatus in

1 the distal part (Fig. 6 a). In the course of further development, the spermatid nucleus  
2 lengthens and acquires a spiral shape. The main volume of the cytoplasm shifts distally and  
3 eventually becomes the residual cytoplasm, which is discharged into the cyst.

4 At a more advanced stage the spermatids are elongated cells (about 4.5  $\mu\text{m}$  long) with  
5 an elongated spiral nucleus, 3.8  $\mu\text{m}$  long and 0.55  $\mu\text{m}$  in diameter (Fig. 6 b, c). Nuclear  
6 chromatin is considerably condensed. In the proximal part, the nucleus has a small  
7 depression, 0.08  $\mu\text{m}$  deep (Fig. 6 d, e), where the acrosome is situated. The acrosome is  
8 formed as a flattened lenticular structure above the proximal part of the condensed nucleus  
9 of the spermatid (Fig. 6 d, e). At an early stage, it is surrounded by a bundle of  
10 microtubules (Fig. 6 e). The acrosome has its own membrane (Fig. 6 d).

11 A bundle of microtubules stretches along the long axis of the cell, reaches its apical  
12 part and passes along one side of the cell periphery (Fig. 6 f-h). The bundle starts from the  
13 kinetid of the axoneme (Fig. 6 i). The flagellum has a typical eukaryotic axoneme (9 + 2)  
14 immersed in a small (depth about 0.6  $\mu\text{m}$ ) invagination of the cytoplasm (Figs. 2 d; 6 f, i).  
15 The basal part of the axoneme is associated with a basal body, which is situated next to the  
16 accessory centriole (Fig. 6 i).

17 The main volume of the cytoplasm is concentrated in the distal third of the cell. It  
18 contains numerous small mitochondria (Fig. 6 c, g, h). Several cisterns derived from the  
19 Golgi apparatus are located near the nucleus and surround the axoneme. A scheme of  
20 sperm structure in *H. irregularis* is shown in Fig. 7.

## 22 **4. Discussion**

23 In this work, we examined the structure of mature sperm and the process of  
24 spermiogenesis in two poecilosclerid demosponges. The information obtained in our study  
25 can be used to clarify several important points concerning the reproductive biology of  
26 sponges. The development of the spermatozoa of *Crellomima imparidens* and *Hymedesmia*  
27 *irregularis* is interesting in several respects.

### 29 **4.1. Spermiogenesis**

30 The last stages of spermiogenesis occur in the two poecilosclerids species studied here  
31 according to a scheme, which is common for all animals with elongated spermatozoa:  
32 chromatin condenses, the nucleus elongates, the acrosome forms in the apical part of the  
33 sperm, the main volume of the cytoplasm (residual cytoplasm) shifts basally and is later  
34 discharged, mitochondria concentrate in the basal part of the sperm. Finally, a

spermatozoid is formed.

## 4.2. Nucleus elongation

In *Crellomima imparidens* and *Hymedesmia irregularis* we found the bundles of microtubules passing along the nucleus during spermiogenesis. These microtubules start from the basal body of the axoneme and stretch along the entire nucleus, reaching the apical pole of the cell, where the acrosome is formed. Perinuclear microtubules starting from basal body have also been described, e.g., in spermatids of *Cryptochiton stelleri* (Mollusca), where they extend anteriorly and encircle the nucleus forming a manchette (Buekland-Nicks et al., 1990). Two other poecilosclerid sponges that have sperm with an elongated nucleus, *Lycopodina occidentalis* and *Crambe crambe*, also have microtubules arranged along the nucleus (Riesgo et al., 2007a; Riesgo & Maldonado, 2009). However, no microtubules have been found in the spermatids of *Lycopodina hypogea*, which has a long helical nucleus (Ereskovsky et al., in prep.).

In *C. imparidens* the microtubules surround the nucleus and are arranged along its longer axis, similarly to the manchette microtubules in the sperms of some metazoans (Kondo et al., 1988; Buekland-Nicks et al., 1990; Russell et al., 1991; Maretia, 1995). Such manchette-like structures have never been reported in Porifera before. In *H. irregularis*, unlike *C. imparidens*, the microtubules pass along only one side of the cell periphery. A single bundle of microtubules was described in the sperm of passerine birds during spermiogenesis (Kondo et al., 1988).

The manchette of microtubules is thought to be responsible for controlling the elongation and shaping of the spherical sperm nucleus to a highly helical form and also for the twisting and/or condensation of chromatin (Russell-Pinto et al., 1983; Lehti & Sironen 2016). This mechanism was described in different animals, for example, in some gastropods (REF: Buckland-Nicks et al., 1983; Maxwell, 1983; Buckland-Nicks et al., 1990), in oligochaetes (Webster & Richards, 1977; Lehti & Sironen, 2019), in passerine birds (Kondo et al., 1988), and in mammals (Russell et al., 1991). Stretched and helical spermatozoa are well known in different invertebrates such as Tubificidae (Oligochaeta) (Erseus & Ferraguti, 1995), Gastrotricha (Ferraguti & Balsamo, 1995), Chelicerata (Alberti, 1995), cephalopod molluscs (Giménez-Bonafé et al., 2002), etc. In sponges this mechanism has been described in *Lycopodina occidentalis* and *Crambe crambe* (Riesgo et al., 2007a; Riesgo & Maldonado, 2009).

### 4.3. Acrosome

Another remarkable feature of spermiogenesis of *Crellomima imparidens* and *Hymedesmia irregularis* is that both species have an acrosome. Sperm with an acrosome has long been thought to be the prerogative of eumetazoans. In Porifera this structure was described for the first time in homoscleromorph *Oscarella* in 1986 (Baccetti et al., 1986; Gaino et al., 1986). Later it was detected in homoscleromorphs *Pseudocortidium jarrei* and *Cortidium candelabrum* (Boury-Esnault & Jamiesson, 1999; Riesgo et al., 2007b), a calcareous sponge *Sycon calcaravis* (Nakamura et al., 1998), and in poecilosclerid demosponges *Crambe crambe* and *Lycopodina hypogea* (Riesgo & Maldonado, 2009; Ereskovsky et al., in prep). The position of the acrosome in *L. hypogea* is typical of all spermatozoa with a dense elongated nucleus and the flagellum on the opposite side of acrosome. However, in *C. crambe*, which has V-shaped spermatozoa, the acrosome is situated close to the basal body at the functionally anterior cell pole.

Moreover, proacrosomal vesicles originating from the Golgi apparatus were described at the apical pole of cells forming during spermiogenesis in many demosponges (Table 1). At later stages of spermiogenesis, the Golgi cisterns disappear but in *Aplysilla rosea*, *Suberiles massa*, *Spongia officinalis*, *Lycopodina occidentalis*, *Petrosia ficiformis*, *Tedania ignis*, and *Geodia phlegraei* dense proacrosomal vesicles of Golgi origin persist near the nucleus of sperms (Tuzet et al., 1970a; Diaz & Connes, 1980; Gaino et al., 1984; Riesgo et al., 2007a; Maldonado & Riesgo 2009; Vasconcellos et al., 2019; Koutsouveli et al., 2020). Proacrosomal vesicles are known in the spermatozoa of many cnidarians (Hinsch, 1974; Franzen, 1996; Larkman & Carter, 1980). They are thought to be a rudimentary acrosome. Nevertheless, proacrosomal vesicles are often involved in fertilization of sea anemones, in which they form aggregations between the nucleus and the plasma membrane in the apical part of the spermatozoon and can be mistaken for the acrosome (Hinsch & Clark, 1973; Dewel & Clark, 1972).

Microtubular transportation of cell organelles and proacrosomal vesicles is well-known in various animals (Hayden et al., 1983; Kierszenbaum & Tres, 2004; Dunleavy et al., 2019). The assembly of the spermatid acrosome depends on the microtubule organization during spermiogenesis (Moreno et al., 2006). We could not trace the formation of acrosomes from the Golgi apparatus in *Crellomima imparidens* and *Hymedesmia irregularis* in detail, and the mechanism of transportation of proacrosomal vesicles remains unresolved. The microtubules in spermatids in the two sponges in our study reach the apical part of the cell where the acrosome is formed. We hypothesize that

the proacrosomal vesicles may be transported along the microtubules, which project all the way to the tip of the sperm.

Thus, we suppose that the acrosome may have appeared more than once in the evolution of the Porifera. Its origin is associated with the features of insemination and fertilization and with the structure of the vitelline envelope and its penetrability for sperm rather than with the taxonomic affiliation or phylogenetic position of the species.

Acrosomes of sponges are variable in shape. Within Homoscleromorpha, they are cup-like in *Oscarella* (Baccetti et al., 1986; Gaino et al., 1986), lens-like in *Pseudocortidium jarrei* (Boury-Esnault & Jamiesson, 1999), and C-shaped in *Corticium candelabrum* (Riesgo et al., 2007b). In the calcarean sponge *Sycon calcaravis* the acrosome is conical, with an electron-dense axial rod in the center (Nakamura et al., 1998). In poecilisclerid demosponges, this structure is also variable. In *Crambe crambe* the acrosome is electron-dense and conical, with an electron-dense acrosomal rod located between it and the nucleus (Riesgo & Maldonado, 2009); in *Lycopodina hypogea* the acrosome is opaque and cone-shaped (Ereskovsky et al., in prep); in *Crellomima imparidens* the acrosome is dome-shaped (this work), and in *Hymedesmia irregularis* the acrosome is formed as a flattened lenticular structure (this work).

#### 4.4. Flagellum and basal apparatus

As in all investigated demosponges, the flagellum in *Crellomima imparidens* and *Hymedesmia irregularis* is already present in the spermatocyte. The axoneme of the spermatid and the sperm flagellum of both investigated sponges have a typical eukaryotic organization (9 + 2) and are located in a deep tunnel-like cytoplasmic depression (1.1 and 0.6  $\mu\text{m}$  depth, correspondingly). It should be noted that in sponges there is a correlation between the shape of sperm and the depth of the cytoplasmic depression (Table 1). In elongated spermatozoa, the tunnel is usually quite deep (e.g., Riesgo et al., 2007a; Riesgo & Maldonado, 2009; Vasconcellos et al., 2019; Ereskovsky et al., in prep.), while in rounded spermatozoa it is either absent or very shallow (e.g. Diaz & Connes, 1980; Tuzet et al., 1970a; Baccetti et al., 1986; Gaino et al., 1984). However, this tunnel, or cytoplasmic depression, of the axoneme is absent in *Halichondria panicea*, which has long (up to 6  $\mu\text{m}$ ) spermatozoa (Barthel & Detmer, 1990). A deep cytoplasmic depression of the axoneme has been described in elongated sperms of other invertebrates such as the sea pen *Pennatula aculeata* (Eckelbarger et al., 1998) and the polychaetes from the genera *Capitella* and *Capitomastus* (Jamieson & Rose, 1989).

As it is typical of invertebrate sperm, the axoneme of the flagellum in both investigated species arises from the basal body. Near this basal body an additional centriole is situated at the angle in both species. It is interesting that an additional centriole was detected only in two other poecilosclerid species, *Iophon proximum* and *I. piceum* (Vasconcellos et al., 2019), out of the ten investigated in this respect. However, it was described in spermatozoa of some sponges from other orders: Axinellida, Haplosclerida, Spongillida, Suberitida, Dactyoceratida, and Dendroceratida (Table 1) (Tuzet et al., 1970a, b; Vacelet, 1979; Diaz & Connes, 1980; Efremova & Papkovskaya, 1980; Paulus, 1989; Riesgo et al., 2008; Maldonado & Riesgo, 2009).

We found a short rootlet starting from a terminal part of the basal body and extending to contact the nucleus in *C. imparidens*. No such structure was detected in *H. irregularis*. In other sponges sperm axoneme rootlet has been described only in poecilosclerids *C. crambe* and *I. piceum* (Riesgo & Maldonado, 2009; Vasconcellos et al., 2019).

#### 4.5. Phylogeny and Taxonomy

According to the comparative morphological concept, there are three morphological types of metazoan spermatozoa: “primitive spermatozoon”, “derived or modified spermatozoon”, and “aberrant spermatozoon” (Franzen, 1956; Reunov, 2005). As pointed out by Riesgo and Maldonado (2009), the spermatozoa of demosponges can be classified as belonging to two structural types, “primitive” or “derived or modified”, depending on the taxa. At the same time, it is sometimes difficult to attribute sperm of the studied species of Demospongiae to one or another category from this classification. The same sperm often has both “primitive” and “derived or modified” traits.

The sperms of *C. imparidens* and *H. irregularis* have such “derived” features as an elongated shape with an elongated helical nucleus, an apical acrosome and manchette-like microtubules. At the same time, they also possess “primitive” traits such as numerous small mitochondria and a large volume of the cytoplasm.

Nevertheless, spermatozoa of poecilosclerids *L. occidentalis*, *L. hypogea*, *C. crambe*, *C. imparidens* and *H. irregularis* are “modified or derived” rather than “primitive”, while spermatozoa of some other species from this order (*Myxilla incrustans*, *Iophon piceum*, *I. proximum*, *Hemimycale columella*, *Tedania ignis*), as well as those of other demosponges, should be classified as “primitive” (sensu Reunov, 2005). **SCHEMES of Sperms?**

The organization of spermatozoa is considered as a sound taxonomic and phylogenetic character (see Afzelius, 1979; Franzén, 1996; Jamiesson et al., 1995; Boury-Esnault &

Jamiesson, 1999). However, the presence of the acrosome in spermatozoa of Poecilosclerida is at variance with this idea. For instance, *Lycopodina hypogea* (fam. Cladorhizidae), *Crambe crambe* (fam. Crambeidae), *Crellomime imparidens* (fam. Crellidae) and *Hymedesmia irregularis* (fam. Hymedesmiidae) have an acrosome; *Lycopodina occidentalis* (fam. Cladorhizidae) and *Tedania ignis* (fam. Tedaniidae) have proacrosomal vesicles, but other investigated sponges from the same order, *Myxilla incrustans* (fam. Myxillidae), *Iophon piceum*, *I. proximum* (fam. Acarnidae), and *Hemimycale columella* (fam. Hymedesmiidae), have none (Table 1). The acrosome is likely to have originated several times in the course of the evolution. In order to verify this hypothesis, thorough studies of the ultrastructural features of insemination in sponges are necessary.

### Acknowledgments

This work was conducted under the IDB RAS Government basic research program in 2021 № 0088-2021-0009 and the grant from the Federal Scientific Policy of Belgium. We thank the Morphology Service of the Mediterranean Institute of Marine and Terrestrial Biodiversity and Ecology (IMBE) for technical support. The authors are warmly grateful to Dr. Philippe Willenz, Royal Belgian Institute of Natural Sciences, for help and discussions, and to Natalia Lentsman for improving the English.

**Conflict of Interest Statement** The authors declare that they do not have any conflict of interest. All authors have read the text and approved the final version of the manuscript.

### Author contributions

AE collected and fixed the animals, interpret the results, write the manuscript, prepared all figures. DT performed the histology, prepared the serial sections, made some schemes, and photographed the sections.

### Data availability statement

The data that support the findings of this study are available from the corresponding author upon request.

### ORCID

Alexander Ereskovsky <https://orcid.org/0000-0003-1079-7204>

Daria Tokina <https://orcid.org/0000-0003-0995-4532>

### References

- Afzelius, B. A. (1978). Fine Structure of the Garfish Spermatozoon, *Journal of Ultrastructural Researches*, 64, 309–314
- Afzelius, B.A. (1979). Sperm Structure in Relation to Phylogeny in Lower Metazoa. In D.W. Fawcett, & J.M. Bedford (Eds.), *The Spermatozoon* (pp. 243–251). Baltimore: Urban and Schwarzenberg.

- 1 Alberti, G. (1995). Comparative spermatology of Chelicerata: review and  
2 perspective. In: B.G.M. Jamieson & A. J. Justine (Eds.), *Advances in spermatozoan*  
3 *phylogeny and taxonomy* (pp. 203–230). Paris: Mémoires du Muséum National d'Histoire  
4 Naturelle.
- 5 Ba, C.T., Marchand, B. (1995). Spermiogenesis, Spermatozoa and Phyletic Affinities  
6 in the Cestoda. In: B.G.M. Jamieson & A. J. Justine (Eds.), *Advances in spermatozoan*  
7 *phylogeny and taxonomy* (pp. 87-95). Paris: Mémoires du Muséum National d'Histoire  
8 Naturelle.
- 9 Baccetti, B. (1984). Evolution of the Spermatozoon. *Italian Journal of Zoology*,  
10 51(1), 25–33
- 11 Baccetti, B., Gaino, E., & Sarà, M. (1986). A sponge with acrosome: *Oscarella*  
12 *lobularis*. *Journal of Ultrastructure and molecular Research*, 94, 195-198
- 13 Barthel, D., & Detmer, A. (1990). The spermatogenesis of *Halichondria panicea*  
14 (Porifera, Demospongiae). *Zoomorphology*, 110, 9-15
- 15 Berruti, G., Ferraguti, M., & Donin, C.L.L. (1978) The aflagellate Spermatozoon of  
16 Ophryotrocha: a line of evolution of fertilization among Polychaetes, *Gamete Gesearch*, 1,  
17 287–292
- 18 Boury-Esnault, N., & Jamieson, B.G.M. (1999). Porifera. In: K.G. Adiyodi & R.G.  
19 Adiyodi (Eds.), *Reproductive Biology of Invertebrates. 9: Pt A Progress in male gamete*  
20 *biology* (pp. 1-41). Oxford & IBH Publishing Co. PVT. LTD, New Delhi, Calcutta
- 21 Buckland-Nicks, J.A, Williams, D., Chia, F.S., & Fontaine, A. (1983). Studies of the  
22 polymorphic spermatozoa of a marine snail. 2. Genesis of the eupyrene sperm. *Gamete*  
23 *Researches*, 7, 19-37
- 24 Buckland-Nicks, J., Koss, R. & Chia, F.-S. (1990). Spermiogenesis in  
25 Polyplacophora, with special reference to acrosome formation. *Zoomorphology*, 109, 179-  
26 188
- 27 Buckland-Nicks, J. (2006). Fertilization in chitons: Morphological clues to  
28 phylogeny. *Venus*, 65, 51-69
- 29 De Vos, L., Rutzler, K., Boury-Esnault, N., Donadey, C., & Vacelet, J. (1991). Atlas  
30 de morphologie des Eponges - Atlas of sponge morphology. Smithsonian Institution Press.  
31 Smithsonian Institution Press. Washington p. 117
- 32 Dewel, W.C., & Clark, W.H. (1972). An ultrastructural investigation of  
33 spermiogenesis and the mature sperm in the Anthozoan *Bunodosoma cavernata* (Cnidaria)  
34 *Journal of Ultrastructure Research*, 40, 417-431
- 35 Diaz, J.P., & Connes, R. (1980). Etude ultrastructural de la spermatogenèse d'une  
36 Demosponge. *Biologie cellulaire*, 38, 225-230
- 37 Dunleavy, L.E.M., O'Bryan, M.K., Stanton, P.G., & O'Donnell, L. (2019). The  
38 cytoskeleton in spermatogenesis. *Reproduction*, 157, R53–R72
- 39 Eckelbarger, K.J., Tyler, P.A., & Langton, R.W. (1998). Gonadal morphology and  
40 gametogenesis in the sea pen *Pennatula aculeata* (Anthozoa: Pennatulacea) from the Gulf  
41 of Maine. *Marine Biology*, 132, 677-690
- 42 Efremova, S.M., & Papkovskaya, M.V. (1980). Spermatogenesis of the sponge  
43 *Lubomirskia baikalensis* (Pallas) ultrastructural investigation. *Archiv of Anatomy,*  
44 *Histology and Embryology*, 79, 88-95
- 45 Efremova, S.M., Ereskovsky, A.V., & Tokina, D.B. (1987). Gametogenesis in  
46 sponges of the family Myxillidae from the White Sea. 2. Spermatogenesis in *Myxilla*  
47 *incrustans* and *Iophon piceus* (Demospongia, Poecilosclerida). *Ontogenez*, 3, 263-268
- 48 Ereskovsky, A.V. (2005) Comparative embryology of Sponges (Porifera). Saint-  
49 Petersburg Univ. Press, Saint-Petersburg (in Russian)

- 1 Ereskovsky, A.V. (2010). The Comparative Embryology of Sponges. Springer-  
2 Verlag, Dordrecht Heidelberg London New York p. 328
- 3 Erseus, C. & Ferraguti M. (1995). The use of spermatozoal Ultrastructure in  
4 phylogenetic studies of Tubincidae. In: B. G. M Jamieson, J. Ausio & J. L. Justine (Eds.),  
5 *Advances in Spermatozoa Phylogeny and Taxonomy*. (pp. 189-201) Paris: Mémoires du  
6 Muséum National d'Histoire Naturelle.
- 7 Ferraguti, M., & Balsamo, M. (1995). Comparative spermatology of Gastrotricha. In:  
8 In : B. G. M Jamieson, J. Ausio & J. L. Justine (Eds.), *Advances in Spermatozoa*  
9 *Phylogeny and Taxonomy*. (pp. 105–117) Paris: Mémoires du Muséum National d'Histoire  
10 Naturelle.
- 11 Franzén, A. (1956). On spermiogenesis, morphology of the spermatozoon, and  
12 biology of fertilization among invertebrates. *Zool Bidrag Uppsala*, 31, 356-482
- 13 Franzén, A. (1996). Ultrastructure of spermatozoa and spermiogenesis in the  
14 hydrozoan *Cordylophora caspia*, with comments on structure and evolution of the sperm in  
15 the Cnidaria and the Porifera. *Invertebrate Reproduction and Development*, 29, 19–26
- 16 Gaino, E., Burlando, B., Zunino, L., Pansini, M., & Buffa, P. (1984). Origin of male  
17 gametes from choanocytes in *Spongia officinalis* (Porifera, Demospongiae). *International*  
18 *Journal of Invertebrate Reproduction and Development*, 7, 83-93
- 19 Gaino, E., Burlando, B., Buffa, P., & Sarà, M. (1986). Ultrastructural study of  
20 spermatogenesis in *Oscarella lobularis* (Porifera, Demospongiae). *International Journal of*  
21 *Invertebrate Reproduction and Development*, 10, 297-305
- 22 Gaino, E., Lepore, E., Reborà, M., Mercurio, M., & Sciscioli, M. (2007). Some steps  
23 of spermatogenesis in *Halichondria semitubulosa* (Demospongiae, Halichondriidae).  
24 *Italian Journal of Zoology*, 74, 117-122
- 25 Giménez-Bonafé, P., Ribes, E., Sautière, P., González, A., Kasinsky, H.E., Kouach,  
26 M., Sautière, P-E., Ausió, J., & Chiva, M. (2002). Chromatin condensation, cysteine-rich  
27 protamine, and establishment of disulphide interprotamine bonds during spermiogenesis of  
28 *Eledone cirrhosa* (Cephalopoda). *European Journal of Cell Biology*, 81, 341–349
- 29 Hayden, J.H., Allen, R.D., & Goldman, R.D. (1983). Cytoplasmic transport in  
30 keratocytes: direct visualization of particle translocation along microtubules. *Cell Motility*,  
31 3, 1-19
- 32 Hinsch, G.H. (1974). Comparative Ultrastructure of Cnidarian Sperm. *American*  
33 *Zoologist*, 14, 457-465
- 34 Hinsch, G.W. & Clark, W.H.J. (1973). Comparative Fine Structure of Cnidaria  
35 Spermatozoa. *Biology of Reproduction*, 8(1), 62–73
- 36 Jamieson, B.G.M., Ausio, J., & Justine J.L. Eds. (1995). Advances in Spermatozoal  
37 Phylogeny and Taxonomy. Mémoires du Muséum national d'Histoire naturelle, p.  
38 564
- 39 Jamieson, B. G., & Rouse, G. W. (1989). The spermatozoa of the polychaeta  
40 (Annelida): an ultrastructural review. *Biological Reviews*, 64, 93-157
- 41 Kaye, H.R., & Reiswig, H.M. (1991). Sexual reproduction in four Caribbean  
42 commercial sponges. I. Reproductive cycles and spermatogenesis. *Invertebrate*  
43 *Reproduction and Development*, 19, 1-11
- 44 Kierszenbaum, A.L., & Tres, L.L. (2004). The acrosome-acroplaxome-manchette  
45 complex and the shaping of the spermatid head. *Archives of histology and cytology*, 67,  
46 271-284
- 47 Kondo, T., Hasegawa, K., & Uchida, T.A. (1988). Formation of the Microtubule  
48 Bundle and Helical Shaping of the Spermatid in the Common Finch, *Lonchura striata* var.  
49 *domestica*. *Journal of ultrastructure and molecular structure research*, 98, 158-168
- 50 Koutsouveli, V., Cárdenas, P., Conejero, M., Rapp, H. T., & Riesgo, A. (2020).

Reproductive Biology of *Geodia* Species (Porifera, Tetractinellida) From Boreo-Arctic North-Atlantic Deep-Sea Sponge Grounds. *Frontiers in Marine Science*, 7, 1–22, doi:10.3389/fmars.2020.595267

Larkman, A.U., & Carter, M.A. (1980). The Spermatozoon of *Actinia equina* L. var. *Mesembryanthemum*. *Journal of the Marine Biological Association of the United Kingdom*, 60, 193–204

Lehti, M.S., & Sironen, A. (2019). Formation and function of the manchette and flagellum during spermatogenesis. *Reproduction*, 151, R43–R54

Maldonado, M., & Riesgo, A. (2009). Gametogenesis, embryogenesis, and larval features of the oviparous sponge *Petrosia ficiformis* (Haplosclerida, Demospongiae). *Marine Biology*, 156, 2181–2197

Maretia, M. (1995). Formation and role of the manchette microtubules in the poultry spermatids. *Acta Veterinaria Brno*, 64, 23–29

Moreno, R.D., Palomino, J., & Schatten, G. (2006). Assembly of spermatid acrosome depends on microtubule organization during mammalian spermiogenesis. *Developmental Biology*, 293, 218–227

Maxwell, W.L. (1983). Mollusca. In: K.G., Adiyodi & R.G., Adiyodi (Eds.), *Reproductive biology of invertebrates, Vol II (Spermatogenesis and sperm function)* (pp. 275 – 319), Wiley, Chichester

Nakamura, Y., Okada, K., & Watanabe, Y. (1998). The ultrastructure of spermatozoa and its ultrastructural change in the choanocyte of *Sycon calcaravis* Hozawa. In: Y. Watanabe & N. Fusetani (Eds.), *Sponge Sciences. Multidisciplinary perspectives* (pp. 179–191). Springer-Verlag, Tokyo

Paulus, W. (1989). Ultrastructural investigation of spermatogenesis in *Spongilla lacustris* and *Ephydatia fluviatilis* (Porifera, Spongillidae). *Zoomorphology*, 109, 123–130

Paulus, W., & Weissenfels, N. (1986). The spermatogenesis of *Ephydatia fluviatilis* (Porifera). *Zoomorphology*, 106, 155–162

Pérez-Porro, A. R., González, J., Uriz, M. J. (2012). Reproductive traits explain contrasting ecological features in sponges: the sympatric poecilosclerids *Hemimyscale columella* and *Crella elegans* as examples. *Hydrobiologia*, 687, 315–330

Reunov, A. (2005). Problem of terminology in characteristics of of Metazoa. *Russian Journal of Developmental Biology*, 36, 335–351

Riesgo, A., & Maldonado, M. (2009). An unexpectedly sophisticated, V-shaped spermatozoon in Demospongiae (Porifera): reproductive and evolutionary implications. *Biological Journal of the Linnean Society*, 97, 413–426

Riesgo, A., Taylor, C., & Leys, S.P. (2007a). Reproduction in a carnivorous sponge: the significance of the absence of an aquiferous system to the sponge body plan. *Evolution & Development*, 9, 618–631

Riesgo, A., Maldonado, M., & Durfort, M. (2007b). Dynamics of gametogenesis, embryogenesis, and larval release in a Mediterranean homosclerophorid demosponge. *Marine and Freshwater Research*, 58, 398–417

Riesgo, A., Maldonado, M., Durfort, M. 2008. Occurrence of somatic cells within the spermatid cysts of demosponges: A discussion of their role. *Tissue and Cell*, 40, 387–396

Russell-Pinto, F., Azevedo, C., & Barandela, T. (1983). Fine structure of the spermatozoa of *Chiton marginatus* (Mollusca: Amphineura), with special reference to nucleus maturation. *Gamete Researches*, 8, 345–355

Russell, L.D., Russell, J.A., Macgregor, G.R., & Meistrich, M.L. (1991). Linkage of Manchette Microtubules to the Nuclear Envelope and Observations of the Role of the Manchette in Nuclear Shaping During Spermiogenesis in Rodents. *The American Journal of Anatomy*, 192, 97–120

- 1 Sukhodolskaya, A. N., & Papkovskaya, M. V. (1985). Electron microscopical study  
2 of spermatogenesis in the fresh-water sponges, *Ephydatia fluviatilis* and *Spongilla*  
3 *lacustris*. *Tsitologiya*, 27, 297-302
- 4 Tripepi, S., Longo, O. M., & La Camera, R. (1984). A new pattern of spermiogenesis  
5 on the sponge *Crambe crambe*: preliminary observations in electron microscopy. Csanady,  
6 A., Rholich, P., & Szabo, D. (Eds.), *Eight European Congress on Electron Microscopy*  
7 (pp. 2073-2074), Budapest
- 8 Tuzet, O., Garrone, R., & Pavans de Ceccatty, M. (1970a). Observations  
9 ultrastructurales sur la spermatogenèse chez la demosponge *Aplysilla rosea* Sch.  
10 (Dendroceratide): une métaplasie exemplaire. *Annales des Sciences naturelles*, 12, 27-50
- 11 Tuzet, O., Garrone, R., & Pavans de Ceccatty, M. (1970b). Origine choanocytaire de  
12 la lignée germinale mâle chez la demosponge *Aplysilla rosea* Schulze (Dendroceratides).  
13 *Compte-Rendu de l'Académie des Sciences de Paris*, 270, 955-957
- 14 Usher, K.M., Sutton, D.C., Toze, S., Kuoc, J., & Fromon, J. (2004). Sexual  
15 reproduction in *Chondrilla australiensis* (Porifera: Demospongiae). *Marine and*  
16 *Freshwater Research*, 55, 123-134
- 17 Vacelet, J. (1979). Quelques stades de la reproduction sexuee d'une éponge  
18 Sphinctozoaire actuelle. In : C. Lévi, & N. Boury-Esnault, (Eds.), *Biologie des Spongiaires*  
19 (pp. 95-101). Paris, Editions du C.N.R.S.
- 20 Vasconcellos, V., Willenz, P., Ereskovsky, A., & Lanna, E. (2019). Comparative  
21 ultrastructure of the spermatogenesis of three species of Poecilosclerida (Porifera,  
22 Demospongiae). *Zoomorphology*, 138(1), 1–12
- 23 Webster, P.M., & Richards, K.S. (1977). Spermiogenesis in the Enchytraeid  
24 *Lumbricillus rivalis* (Oligochaeta: Annelida). *Journal of Ultrastructure Research*, 61, 62-  
25 77
- 26 Watson, N.A., & Rhode, K. (1995). Sperm and spermiogenesis of the “Turbellaria”  
27 and implications for the phylogeny of the phylum Platyhelminthes. In: B. G. M Jamieson,  
28 J. Ausio & J. L. Justine (Eds.), *Advances in Spermatozoa Phylogeny and Taxonomy* (pp.  
29 37-54) Paris: Mémoires du Muséum National d'Histoire Naturelle.
- 30 Willenz P., Hartman W.D., 2004. Storage cells and spermatocysts in the Caribbean  
31 *Coralline sponge* *Goreauia auriculata* (Astroscleridae, Agelasida, Demospongiae): a  
32 relationship? *Boll. Mus. Ist. Biol. Univ. Genova*, 68, 673-681

## Figure legends

**Figure 1.** *In situ* photographs of the investigated poecilosclerids. (a) *Crellomima imparidens*; (b) *Hymedesmia irregularis*. ca – aquiferous system canals; o – oscula.

**Figure 2.** TEM micrographs of spermatid cysts of investigated species. (a) The cyst with early spermatids in *Crellomima imparidens*. (b) The cyst in *Hymedesmia irregularis*. (c) The cyst with later spermatids and spermatozooids in *C. imparidens*. (d) The cyst with later spermatids in *H. irregularis*. co – cytoplasm outgrowths of follicular cells; fc – follicular cell; n – nucleus; ph – phagosomes; st – spermatids. Scale bars: a-c – 5 µm; d – 2 µm.

**Figure 3.** TEM images of spermiogenesis in *Crellomima imparidens*. (a) Early spermatids in the cyst. (b) Spermatids connected through cytoplasmic bridges (br). (c,d) Longitudinal section of the spermatids at the later stages during cells stretching and the nuclei twisting. (e) Longitudinal section of the basal part of sperm, showing the microtubules (arrowheads) begin from the basal body of the axoneme and are uniformly extending along the nucleus. (f, g) Cross-section of the sperms at the basal (f) and central (g) parts showing the microtubules (arrowheads) manchette surrounding the nucleus. (h–j) Longitudinal sections of the apical part of sperms with the acrosome. (k, l) Cross-sections of the acrosome. (m) Cytoplasm depression at the base of sperm flagellum. (n, o) Kinetid of sperm axoneme with basal body (bb), accessory centriole (ac) and the short rootlet (arrowhead). a – acrosome; ac – accessory centriole; bb – basal body of sperm axoneme; f – flagellum; n – nucleus; rc – residual cytoplasm. Scale bars: a – 5 µm; b – 2 µm; c–e – 1 µm; f – g – 0.5 µm; h–l – 0.5 µm; m – 2 µm; n – 1 µm; o – 2 µm.

**Figure 4.** Longitudinal (a) and corresponding cross-sections (b–f) of sperm in *Crellomima imparidens*. a – acrosome; ac – accessory centriole; bb – basal body of sperm axoneme; mt – microtubules; mtc – mitochondria; n – nucleus. Scale bars: a, b – 0.5 µm.

**Figure 5.** Schematic drawing of *Crellomima imparidens* spermatozoon. (a) Longitudinal section. (b) Cross and longitudinal section of the apical part of sperm. a – acrosome; ac – accessory centriole; bb – basal body of sperm axoneme; f – flagellum; mt – microtubules; mtc – mitochondria; n – nucleus.

**Figure 6.** Spermiogenesis in *Hymedesmia irregularis*. (a) Early spermatids of *Hymedesmia irregularis*. (b, c) Spermatids at the late stages with an elongated spiral-bound nucleus (c). (d, e) Apical part of the nuclei in late spermatids with small depression (arrowheads) and an acrosome. (f–h) Longitudinal (f, g) and cross-sections of spermatids, showing the bundles of microtubules stretches along the long axis of the cell. (i) Kinetid of sperm axoneme with basal body (bb), accessory centriole (ac); the microtubule bundle starts from the axoneme's kinetid (arrowheads). a – acrosome; f – flagellum; mt – microtubules; n – nucleus rc – residual cytoplasm; st – spermatids. Scale bars: a, b, c, f, g, h, i – 1 µm; d, e – 0.5 µm.

**Figure 7.** Schematic drawing of *Hymedesmia irregularis* spermatozoon. (a) Longitudinal section. (b) Cross and longitudinal section of the apical part of sperm. a – acrosome; ac – accessory centriole; bb – basal body of sperm axoneme; f – flagellum; mt – microtubules; mtc – mitochondria; n – nucleus.