



HAL
open science

ROCHECHOUART 2017-DRILLING CAMPAIGN: FIRST RESULTS

P Lambert, C Alwmark, D Baratoux, S Bouley, A Brack, P Bruneton, E Buchner, P Claeys, M Dence, A Nomade, et al.

► **To cite this version:**

P Lambert, C Alwmark, D Baratoux, S Bouley, A Brack, et al.. ROCHECHOUART 2017-DRILLING CAMPAIGN: FIRST RESULTS. 49th Lunar and Planetary Science Conference, Mar 2018, The Woodlands, Texas, United States. hal-03543017

HAL Id: hal-03543017

<https://hal-cnrs.archives-ouvertes.fr/hal-03543017>

Submitted on 25 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ROCHECHOUART 2017-DRILLING CAMPAIGN: FIRST RESULTS. P. Lambert¹, C. Alwmark², D. Baratoux³, S. Bouley⁴, A. Brack⁵, P. Bruneton⁶, E. Buchner⁷, P. Claeys⁸, M.R. Dence⁹, A.Courtin Nomade¹⁰, I. Duhamel Achin¹¹, J.P. Floch¹², B.M. French¹³, C. Fudge¹⁴, J. Gattacceca¹⁵, R.L. Gibson¹⁶, S. Goderis¹⁷, R.A.F. Grieve¹⁸, K.V. Hodges¹⁹, F. Hörz²⁰, M. Humayun²¹, F. Jourdan²², S.P. Kelley²³, T. Kenkmann²⁴, D.A. Kring²⁵, F. Langenhorst²⁶, M.R. Lee²⁷, P. Lindgren²⁸, J. Lofi²⁹, J.P. Lorand³⁰, B. Luais³¹, V. Masaitis³², A. Meunier³³, C.B. Moore³⁴, J. Ormö³⁵, G.R. Osinski³⁶, S. Petit³⁷, P.A. Pezard³⁸, M. Poelchau³⁹, J. Pohl⁴⁰, Y. Quesnel⁴¹, C. Ramboz⁴², H. Reeves⁴³, P. Rochette⁴⁴, H.M. Sapers⁴⁵, M. Schmieder⁴⁶, P.H. Schultz⁴⁷, S.P. Schwenzer⁴⁸, T. Sharp⁴⁹, C.S. Shoemaker⁵⁰, S.L. Simpson⁵¹, D. Stöffler⁵², E. Sturkell⁵³, H. Trumel⁵⁴, E. Walton⁵⁵, F. Westall⁵⁶, A. Wittmann⁵⁷, and K. Wünnemann⁵⁸.

¹CIRIR-Center for International Research on Impacts and on Rochechouart-87600 Rochechouart-France, lambertbdx@gmail.com, ²carl.alwmark@geol.lu.se, ³david.baratoux@get.obs-mip.fr, ⁴sylvain.bouley@u-psud.fr, ⁵brack@cnsr-orleans.fr, ⁶p.bruneton@orange.fr, ⁷Elmar.Buchner@hs-neu-ulm.de, ⁸phclaeys@vub.ac.be, ⁹alexandra.courtin@unilim.fr, ¹⁰mrdenca@rogers.com, ¹¹i.duhamelachin@brgm.fr, ¹²jpfloch@gmail.com, ¹³FRENCHB@si.edu, ¹⁴cfudge@asu.edu, ¹⁵gattacceca@cerege.fr, ¹⁶roger.gibson@wits.ac.za, ¹⁷Steven.Goderis@vub.ac.be, ¹⁸rgrieve@nrcan.gc.ca, ¹⁹kvhodges@asu.edu, ²⁰fhorz@aol.com, ²¹humayun@magnet.fsu.edu, ²²F.Jourdan@exchange.curtin.edu.au, ²³simon.kelley@open.ac.uk, ²⁴thomas.kenkmann@geologie.uni-freiburg.de, ²⁵Kring@lpi.usra.edu, ²⁶falko.langenhorst@uni-jena.de, ²⁷Martin.Lee@glasgow.ac.uk, ²⁸Paula.Lindgren@glasgow.ac.uk, ²⁹johanna.lofi@gm.univ-montp2.fr, ³⁰Jean-Pierre.Lorand@univ-nantes.fr, ³¹luais@crpg.cnrs-nancy.fr, ³²vcmts@mail.ru, ³³alain.meunier@univ-poitiers.fr, ³⁴CMoore@asu.edu, ³⁵ormoj@cab.inta-csic.es, ³⁶gosinski@uwo.ca, ³⁷sabine.petit@univ-poitiers.fr, ³⁸ppezard@gulliver.fr, ³⁹michael.poelchau@geologie.uni-freiburg.de, ⁴⁰pohl@geophysik.uni-muenchen.de, ⁴¹quesnel@cerege.fr, ⁴²cramboz@cnsr-orleans.fr, ⁴³nelly@hubertreeves.info, ⁴⁴rochette@cerege.fr, ⁴⁵haley.sapers@gmail.com, ⁴⁶martin@suevite.com, ⁴⁷peter_schultz@brown.edu, ⁴⁸susanne.schwenzer@open.ac.uk, ⁴⁹tom.sharp@asu.edu, ⁵⁰mrscomet3@aspect1.net, ⁵¹ssimps56@uwo.ca, ⁵²dieter.stoeffler@mfn-berlin.de, ⁵³erik.sturkell@gvc.gu.se, ⁵⁴herve.trumel@wanadoo.fr, ⁵⁵ewalton@ualberta.ca, ⁵⁶frances.westall@cnsr-orleans.fr, ⁵⁷axel.wittmann@asu.edu, ⁵⁸kai.wuennemann@mfn-berlin.de

We report on the first comprehensive scientific drilling campaign to be completed in the Rochechouart impact structure, France (Fig. 1), describing the various holes and the first visual inspection of the major lithologies encountered (Fig. 2), in reference to the IUGS nomenclature [1].

Results: 18 holes have been drilled vertically, ranging from 1 to 120 m in depth. The holes are located at 8 sites spread along two 10 km-long radial traverses across the center of the structure (Fig. 1).




Fig.1. Drill sites and drillholes characteristics. Regional map after [2], and geological data after [3].

The cumulative depth drilled is 515 m, with 99.7% recovery. The final numbers may slightly change following forthcoming geophysical logging, to be performed early 2018 by the CNRS (Geosciences Montpellier) with the same rationale, objectives and down-

hole sensors they used for the IODP Chicxulub Expedition 364 [4].

In the allochthonous units, maximum clast size is < 0.5 m and rarely exceeds the core diameter (86 mm). The two shallow holes at site 1 mainly sample vertical flow banded, fine grained impactoclastite dikes similar to those described in [3]. The 120 m SC2 hole located near the top of the Chassenon breccia deposits (site 1, Fig. 1) intersects i) "suevite" (impact melt-bearing breccia) down to 88 m, ii) monomict breccia down to 113 m with clasts comprising fine grained and strongly fractured gneiss, and iii) "massive" locally cataclased gneiss cut by thin monomict breccia dykes. The contact at the bottom of the suevite is transitional. Below the contact, the matrix of the breccia and fractures in the clasts are more or less systematically healed, mainly by hydrothermal carbonates, whereas above the contact, hydrothermal effects appear to be restricted to the melt clasts. The suevite clearly displays two different units. The upper unit fits the classical description of "suevite" according to [1]; comparatively melt clasts in the lower unit are rare. The transition occurs at a depth of ~40 m. Above, clasts are randomly oriented and melt fragments are usually equant. However, clast size distribution and clast/matrix proportion are not random. Unlike the upper suevite, the lower melt-bearing unit locally shows oriented textures and evidence of flow. Dilatancy effects are suspected from the occurrence of

“cloudy” monomict breccia lenses interpreted as “popped out” clasts in the lower suevite. The underlying monomict breccia unit is cut by multiple horizontal pseudotachylitic breccia dikes. Several cm- to sub-meter-wide impactoclastite dikes cut the upper suevite. They are not observed below 30 m.

Holes SC4, 5 and 6 (site 2, Fig.1) intersecting the bottom of the Chassenon deposits resemble the bottom of the SC3 hole located 1 km from site 2.

The 60 m deep holes SC7, 11, 16 and 17 (circled zone on Fig. 1), start with massive, heterogeneous and textured clast-rich impact melt rocks. Melt-in-melt textures are abundant. The units become more homogeneous with depth without homogenizing completely. All impact melts display evidence of flow. The units extend respectively 15, 12, 10 and 20 m below surface. SC7 (site 3, Fig. 1) is distinguished by vesicular clasts at the top of the unit. SC17 (site 7, Fig.1) displays a 2 m thick, massive, yellow impact melt rock intercalated within a purple-red impact melt rock unit. The latter bears small (mm) vesicles and wormlike vesicles. It lies directly on highly deformed crystalline basement rocks. In SC7, SC11 and SC16, the impact melt rock units merge into suevites and “melt-poor suevites”, with sharp variations in texture, clast size and clast shape, down to 20, 40 and 17 m, respectively, where basement rocks are intersected. The latter consist of granitoid and granitoid gneiss displaying cataclastic texture and anastomosed veining. They are locally intersected by a variety of breccia dikes (monomict, polymict lithic, impact melt and pseudotachylitic breccias, and pseudotachylite). Hydrothermal healing is less developed than in the external cores.

SC14 (site 5, Fig.1) is located at the foot of the 40 m thick historical Rochechouart cliff exposing polymict lithic breccia. The hole starts with 1.5 m of altered polymict breccia bearing some rounded clasts and flow-like features in the matrix. It merges into a matrix-rich flow-foliated monomict breccia with rounded clasts. At 4 m depth it is cut by a 1 m thick pseudotachylitic breccia, below which occurs a highly cataclased granitic gneiss that is locally brecciated and cut by a series of breccia dikes (polymict lithic, pseudotachylitic and banded melt-rock). Fractures filled by hydrothermal minerals cross-cut all the units and become more abundant at ~30 m depth and below.

SC15 (site 5, Fig.1) samples 1 m of yellow clast-rich impact melt rock.

SC18 (site 8, Fig. 1), located at the top of the Champagnac quarry, starts with 12 m of monomict gneissic breccia with rounded clasts and varied proportion of matrix. From 10 to 12 m, the proportion of matrix increases and, the size and angularity of clasts decreases. At 12 m, it merges into a complex 2 m thick

unit of pseudotachylitic breccia alternating with white and black altered mylonite or pseudotachylite, underlain by 2 m of brecciated gneiss. An ~40 cm thick mafic dike with black veins marks the boundary with migmatitic rocks, gneiss, granitoid and granodiorite, that are all highly cataclased, sheared and veined to 60 m depth. This unit is cut by monomict, polymict lithic, and pseudotachylitic, breccia dikes. All units display multiple stages of veining, fracturing and hydrothermal healing.

Conclusions and Perspectives: The quantity and quality of materials recovered by the drilling campaign has exceeded expectations [5] enabling CIRIR and PIs to fully develop their projects (list on cirir-edu.org). Core samples will be assigned following the first sample EOI party scheduled for late Spring 2018. CIRIR’s second call for PI projects is now open until late 2018 (see website for details).

Acknowledgments: Special thanks to P. Poupart and to the Reserve Naturelle Nationale de l’Astroblème de Rochechouart-Chassenon responsible for the drillings realized in the sites of the Reserve.

References: [1] Stöffler, D., and R. A. F. Grieve (2007) *Cambridge University Press*, 82-92. [2] Bobée C. et al. (2010) *Near Surface Geophysics* 8, 259-270. [3] Lambert P. (2010) *GSA Spec Pap.* 465, 505–541. . [4] Morgan J. V. et al. (2017) *Science*, 354, 878–882. [5] Lambert P. et al. (2016) MAPS, Abstract, #6471.pdf.




Fig.2. Representative impactite lithologies and features of the CIRIR cores (see each core image for provisional description and location).