



HAL
open science

Self-consistent Monte Carlo model of particle acceleration by relativistic shocks

S Osipov, A Bykov, Martin Lemoine

► **To cite this version:**

S Osipov, A Bykov, Martin Lemoine. Self-consistent Monte Carlo model of particle acceleration by relativistic shocks. *Journal of Physics: Conference Series*, IOP Publishing, 2021, 2103 (1), pp.012014. 10.1088/1742-6596/2103/1/012014 . hal-03796196

HAL Id: hal-03796196

<https://hal-cnrs.archives-ouvertes.fr/hal-03796196>

Submitted on 5 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License

PAPER • OPEN ACCESS


Self-consistent Monte Carlo model of particle acceleration by relativistic shocks

To cite this article: S M Osipov *et al* 2021 *J. Phys.: Conf. Ser.* **2103** 012014

View the [article online](#) for updates and enhancements.

You may also like

- [TESTING IN SITU ASSEMBLY WITH THE KEPLER PLANET CANDIDATE SAMPLE](#)
Brad M. S. Hansen and Norm Murray
- [Influence of the angular scattering of electrons on the runaway threshold in air](#)
O Chanrion, Z Bonaventura, A Bourdon et al.
- [Evolution of Surface Roughness during Copper Electrodeposition in the Presence of Additives: Comparison of Experiments and Monte Carlo Simulations](#)
Timothy O. Drews, Jason C. Ganley and Richard C. Alkire



ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Presenting more than 2,400 technical abstracts in 50 symposia

Register now!

ECS Plenary Lecture featuring M. Stanley Whittingham,
Binghamton University
Nobel Laureate –
2019 Nobel Prize in Chemistry

The banner features a portrait of M. Stanley Whittingham, a Nobel medal, and a background image of a person interacting with a futuristic interface.

Self-consistent Monte Carlo model of particle acceleration by relativistic shocks

S M Osipov¹, A M Bykov^{1,2}, M Lemoine³

¹ Ioffe Institute, 26 Politekhnicheskaya st., St. Petersburg 194021, Russia

² Peter The Great St. Petersburg Polytechnic University, 29 Politekhnicheskaya st., St. Petersburg 195251, Russia

³ Institut d'Astrophysique de Paris, CNRS - Sorbonne Universite, 98 bis boulevard Arago, F-75014 Paris, France

E-mail: osm2004@mail.ru

Abstract. We present a self-consistent Monte Carlo model of particle acceleration by relativistic shock waves. The model includes the magnetic field amplification in the shock upstream by cosmic ray driven plasma instabilities. The parameters of the Monte Carlo model are obtained based on PIC calculations. We present the spectra of accelerated particles simulated in the frame of the model.

1. Introduction

The processes of particle acceleration and the magnetic field amplification by shocks that occur when a powerful relativistic current collides with the environment are appropriate for many astrophysical objects. Such flows are formed, for example, in the propagation of a pulsar wind, the explosion of relativistic supernovae, and the formation of gamma-ray bursts. These objects are powerful sources of non-thermal radiation. Relativistic shock waves in astrophysical objects are often collisionless. The particles are accelerated to high energies by such shock waves according to the first-order Fermi mechanism, when they are scattered by magnetic fluctuations. The distribution function of accelerated particles has a significant anisotropy in the upstream of a relativistic collisionless shock wave. This anisotropy leads to the development of plasma instabilities that amplify magnetic fields [1–3]. In the same way, the pressure of accelerated particles modifies the prefront of the shock wave [4–9]. Thus, the problem of particle acceleration by a relativistic collisionless shock wave is essentially nonlinear, since the propagation of particles is determined by magnetic fields, and the amplification of fields is a function of the distribution of accelerated particles.

We develop a stationary self-consistent Monte Carlo model of particle acceleration by relativistic shock waves, which takes into account the amplification of magnetic fields by plasma instabilities. Model allows us to construct the particle distribution function, plasma flow profiles, and magnetic fluctuation spectra in an iterative scheme that implements the laws of conservation of momentum and energy near the shock wave front. PIC (particle in cell) calculations [10–17] based on the first principles allow us to solve this problem more accurately. PIC calculations have very significant limitations on the size of the calculation space due to the limited computing power. We compare the results of PIC calculations and the results of our Monte Carlo simulation,



which allows us to set the instability parameters necessary for use in the Monte Carlo model. After calibrating the Monte Carlo model with the PIC results, the calculated spatial regions of the Monte Carlo model can be extended to the actual flow scales of the studied astrophysical objects, which are not available for PIC modeling.

2. Description of the Monte Carlo model

The Monte Carlo model of particle acceleration by an unmagnetized relativistic stationary plane-parallel shock presented here is self-consistent. The model in an iterative scheme constructs the shock flow structure which obeys the momentum and energy conservation laws. The iteration scheme in the relativistic Monte Carlo model is similar to that used in the modeling of non-relativistic shocks with the magnetic field amplification developed earlier [18–21]. The plasma in the model is divided between the background and the accelerated particles. The accelerated particles are those that have crossed the shock wave front at least once from the downstream to the upstream. In most non-relativistic calculations, the background plasma is described hydrodynamically in most of the upstream, which makes it possible to speed up greatly the operation of the numerical scheme. In the calculations of this paper, the entire plasma is described as particles in the entire computational domain.

The pressure gradient of the accelerated particles modifies the upstream momentum fluxes of the background plasma. The distribution function of accelerated particles has a significant anisotropy in the upstream, which leads to the development of plasma instabilities. In this paper, we describe the magnetic field amplification in the upstream by the following equation

$$u_w(x) \frac{dW(x)}{dx} + \frac{\alpha_w}{\gamma_w(x)} \frac{d(\gamma_w(x) u_w(x))}{dx} W(x) = \frac{G(x)}{\gamma_w(x)} W(x), \quad (1)$$

where $u_w(x)$ and $\gamma_w(x)$ are velocity and Lorentz factor of the scattering centre rest frame relative to the shock rest frame accordingly, $W(x)$ and $G(x)$ are turbulence energy density and the growth rate of plasma instabilities in the rest frame of scattering centers accordingly. The model assumes that the magnetic field amplification occurs in a narrow range of wave numbers. The second term of the left side of the equation determines the adiabatic amplification of magnetic fluctuations with a change in the velocity of the scattering centers. Thus, the parameter α_w determines the adiabatic amplification of the magnetic field, in the calculations $\alpha_w = 2$. The x coordinate is calculated from the shock front, along the normal to the front. Negative values x correspond to the upstream, positive values of x correspond to the downstream. In the far upstream, there is a free-escape boundary located at $x = x_{FEB}$. Any accelerated particle which is crossing the boundary leaves the simulation domain. Near the free-escape boundary, the Lorentz factors of the background plasma flow and the rest frame of scattering centers coincide and are equal to the Lorentz factor of the shock γ_{sh} .

The scattering of particles occurs in the rest system of the scattering centers and in this system they are isotropic and elastic. The free path length of particles in the scattering center rest frame in this paper

$$\lambda_{mfp}(x) = \frac{r_g^2(x)}{L_c(x)}, \quad (2)$$

where $L_c(x)$ is the turbulence correlation length, which is determined by the range of wave numbers in which the magnetic fields are amplified, the particle gyroradius is

$$r_g(x) = \frac{cp}{eB(x)} \quad (3)$$

where c is the speed of light, e is the particle charge, p is the momentum modulus of a particle in the scattering center rest frame,

$$B(x) = \sqrt{4\pi W(x)}. \quad (4)$$




Figure 1. The energy flux for calculation with $\gamma_{sh} = 10$. The green curve corresponds to the self-consistent calculation. The blue curve corresponds to the unmodified calculation. F_{E0} is the energy flux in the unperturbed far upstream. $r_{g0} = \frac{m_e c^2}{eB_0} \approx 5.7 \cdot 10^8 \text{cm}$.




Figure 2. The particle distribution functions in the shock rest system are shown by solid curves for $x = 0$ (the shock front) and dashed for $x = x_{FEB}$ (the free-escape boundary). The green curves correspond to $\gamma_{sh} = 10$. The blue curves correspond to $\gamma_{sh} = 100$.

The growth rate of plasma instabilities $G(x)$ and the turbulence correlation length $L_c(x)$ were obtained by comparing the simulation results with the results of PIC calculations [15–17].




Figure 3. The magnetic field profile near the shock. The green curves correspond to $\gamma_{sh} = 10$. The blue curves correspond to $\gamma_{sh} = 100$. $r_{g0} = \frac{m_e c^2}{e B_0} \approx 5.7 \cdot 10^8 \text{ cm}$.

3. Results

Based on our model in this paper we perform a simulation of particle acceleration and magnetic field amplification by shocks with Lorentz factors $\gamma_{sh} = 10$ and $\gamma_{sh} = 100$. The electron-positron plasma is considered. The value of the other calculation parameters is the same: $x_{FEB} = 10^3 r_{g0}$, where $r_{g0} = \frac{m_e c^2}{e B_0}$, m_e is the electron mass, $B_0 = 3 \mu G$ is the root-mean-square value of the magnetic field in the far upstream in the background plasma rest system, $n_0 = 4 \cdot 10^{-3} \text{ cm}^{-3}$ is the total concentration of electrons and positrons in the far upstream in the background plasma rest system.

Figure 1 is given to demonstrate the convergence of the iterative scheme of our Monte Carlo model. This figure shows the energy flux near the shock for two cases: a converged nonlinear solution of the problem and an unmodified calculation. The unmodified calculation does not include self-consistent magnetic field amplification and modification of the upstream by the momentum flow of accelerated particles. As can be seen from Figure 1, in the unmodified case, the energy flow near the shock wave is not conserved.

The results of the self-consistent simulation are shown in the Figures 2 and 3.

The particle distribution functions in the shock rest system are shown on Figure 2. As can be seen from this Figure for the values of the particle momentum $\leq \gamma_{sh} m_e c$, the particle distribution function has the form of a thermal spectrum. For the values of the particle momentum $> \gamma_{sh} m_e c$, the particle distribution function has the form of a power spectrum. The spectrum slope in the power region is close to -4.22. This slope of the spectrum is obtained both in calculations for particle acceleration by relativistic shocks and in the test-particle regime. Since the spatial domain of calculations in this paper is an order of magnitude larger than that achievable in PIC modeling the value of the maximum particle momentum is several times greater than in PIC modeling for similar parameters of the problem.

The simulated magnetic field profiles are shown on Figure 3. A sharp increase in the turbulent magnetic field occurs near the free-escape boundary due to Weibel and filamentation plasma instabilities [14, 17, 22–26]. Closer to the shock front, the magnetic field amplification occurs mainly adiabatically due to the second term in the left part of the equation (1). The magnetic

field amplification effect is essential to explain the observed non-thermal synchrotron emission from the shocks driven by relativistic outflows.

Acknowledgments

A.M. Bykov and S.M. Osipov acknowledge support from RSF grant 21-72-20020.

Some results of the work were obtained using computational resources of Peter the Great Saint-Petersburg Polytechnic University Supercomputing Center (<http://scc.spbstu.ru>) and Joint SuperComputer Center of the Russian Academy of Sciences (<http://www.jscs.ru>).

References

- [1] Bell A R 2004 *MNRAS* **353** 550–558
- [2] Zirakashvili V N, Ptuskin V S and Völk H J 2008 *ApJ* **678** 255–261
- [3] Bykov A M, Brandenburg A, Malkov M A and Osipov S M 2013 *Space Sci. Rev.* **178** 201–232
- [4] Blandford R and Eichler D 1987 *Phys. Reports* **154** 1–75
- [5] Berezhko E G and Krymskiĭ G F 1988 *Soviet Physics Uspekhi* **31** 27–51
- [6] Jones F C and Ellison D C 1991 *Space Sci. Rev.* **58** 259–346
- [7] Malkov M A and Drury L O 2001 *Reports of Progress in Physics* **64** 429–481
- [8] Zirakashvili V N and Ptuskin V S 2012 *Astroparticle Physics* **39** 12–21
- [9] Marcowith A, Bret A, Bykov A, Dieckman M E, O’C Drury L, Lembège B, Lemoine M, Morlino G, Murphy G, Pelletier G, Plotnikov I, Reville B, Riquelme M, Sironi L and Stockem Novo A 2016 *Reports on Progress in Physics* **79** 046901
- [10] Spitkovsky A 2008 *ApJ* **682** L5
- [11] Keshet U, Katz B, Spitkovsky A and Waxman E 2009 *ApJ* **693** L127–L130
- [12] Sironi L, Spitkovsky A and Arons J 2013 *ApJ* **771** 54
- [13] Sironi L, Keshet U and Lemoine M 2015 *Space Sci. Rev.* **191** 519–544
- [14] Pelletier G, Gremillet L, Vanthieghem A and Lemoine M 2019 *Phys. Rev. E* **100** 013205
- [15] Lemoine M, Vanthieghem A, Pelletier G and Gremillet L 2019 *Phys. Rev. E* **100** 033209
- [16] Lemoine M, Pelletier G, Vanthieghem A and Gremillet L 2019 *Phys. Rev. E* **100** 033210
- [17] Lemoine M, Gremillet L, Pelletier G and Vanthieghem A 2019 *Phys. Rev. Lett.* **123** 035101
- [18] Ellison D C, Warren D C and Bykov A M 2013 *ApJ* **776** 46
- [19] Bykov A M, Ellison D C, Osipov S M and Vladimirov A E 2014 *ApJ* **789** 137
- [20] Bykov A M, Ellison D C, Marcowith A and Osipov S M 2018 *Space Sci. Rev.* **214** 41
- [21] Osipov S M, Bykov A M and Ellison D C 2019 *Journal of Physics Conference Series* **1400** 022004
- [22] Weibel E S 1959 *Phys. Rev. Lett.* **2** 83–84
- [23] Yoon P H 1989 *Physics of Fluids B* **1** 1336–1338
- [24] Achterberg A and Wiersma J 2007 *A&A* **475** 1–18
- [25] Achterberg A, Wiersma J and Norman C A 2007 *A&A* **475** 19–36
- [26] Califano F, Pegoraro F and Bulanov S V 1997 *Phys. Rev. E* **56** 963–969